




# FIU

FLORIDA INTERNATIONAL UNIVERSITY  
*Miami's public research university*

## **The 2008 Terrestrial Vegetation of Biscayne National Park, FL, USA Derived From Aerial Photography, NDVI, and LiDAR**

Pablo L. Ruiz, Patricia A. Houle, & Michael S. Ross



June 17, 2008

Funded by the National Park Service South Florida / Caribbean Network  
Cooperative Agreement: H5000 06 0104

Southeast Environmental Research Center  
11200 S.W. 8th Street, OE 148 • Miami, FL 33199 • Tel: (305) 348-3095 • Fax: (305) 348-4096 • [www.fiu.edu](http://www.fiu.edu)

# Contents

Contributors: _____	iii
Executive Summary: _____	iv
Introduction: _____	1
Methods: _____	4
Imagery: _____	4
Vegetation Classification System: _____	5
Shoreline Segmentation (Definiens Professional® v5): _____	7
Normalized Difference Vegetation Index (NDVI): _____	9
Light Detection & Ranging (LiDAR): _____	10
Unsupervised Classification: _____	12
Geodatabase Design and Mapping: _____	13
Accuracy Assessment & Ground-truthing: _____	15
Results & Discussion: _____	17
Vegetation Classification System: _____	17
NDVI & LiDAR: _____	20
Unsupervised Classification: _____	21
Vegetation: _____	24
Conclusion: _____	28
Acknowledgements: _____	29
Literature Cited: _____	30
Plate 1: <i>The 2008 Terrestrial Vegetation of Biscayne National Park.</i> _____	32
Appendices:	
Appendix 1: <i>Hierarchical Classification System with Community Descriptions.</i> _____	A1-1
Appendix 2: <i>Definiens Professional® v5 Segmentation Algorithm.</i> _____	A2-1

## **Contributors:**

**Patricia A. Houle**

Florida International University  
International Hurricane Research Center  
11200 S.W. 8<sup>th</sup> Street  
Miami, FL 33199  
houlep@fiu.edu

**Michael S. Ross**

Florida International University  
Southeast Environmental Research Center/Environmental Studies Department  
11200 S.W. 8<sup>th</sup> Street  
Miami, FL 33199  
rossm@fiu.edu

**Pablo L. Ruiz**

Florida International University  
Southeast Environmental Research Center  
11200 S.W. 8<sup>th</sup> Street  
Miami, FL 33199  
ruizp@fiu.edu

## Executive Summary:

Established as a National Park in 1980, Biscayne National Park (BISC) comprises an area of nearly 700 km<sup>2</sup>, of which most is under water. The terrestrial portions of BISC include a coastal strip on the south Florida mainland and a set of Key Largo limestone barrier islands which parallel the mainland several kilometers offshore and define the eastern rim of Biscayne Bay. The upland vegetation component of BISC is embedded within an extensive coastal wetland network, including an archipelago of 42 mangrove-dominated islands with extensive areas of tropical hardwood forests or hammocks. Several databases and vegetation maps describe these terrestrial communities. However, these sources are, for the most part, outdated, incomplete, incompatible, or/and inaccurate. For example, the current, Welch et al. (1999), vegetation map of BISC is nearly 10 years old and represents the conditions of Biscayne National Park shortly after Hurricane Andrew (August 24, 1992). As a result, a new terrestrial vegetation map was commissioned by The National Park Service Inventory and Monitoring Program South Florida / Caribbean Network.

A vector map was developed using: a comprehensive set of 2005 5-band (Red, Green, Blue, NIR, and Pan-Chromatic) 30cm pixel aerial photographs; NDVI (calculated from the 2005 aeriels); 2002 LiDAR data, available only for mainland portions of BISC; and over 1,000 ground reference points. In general, NDVI helped delineate low productivity zones (Mangrove Scrub) and Non-Vegetative features from adjacent Shrubland and Forest communities. The availability of LiDAR for the mainland proved invaluable and greatly enhanced the overall map resolution and accuracy. In conjunction with traditional aerial photo-interpretations, *Definiens Professional*® v5 remote sensing software was used to create 1:300 scale shorelines. However, we failed to derive an algorithm in *Definiens Professional*® v5 capable of consistently and accurately segmenting the varied community types found in this region. Vegetation communities were classified to the highest feasible level of resolution within a six-tiered hierarchical vegetation classification system. Level 3 of the hierarchy was the minimum resolution accepted, but some communities were mapped to Level 6. However, not all communities mapped or observed in the field were described in the original classification system. As a result, we modified the classification system, adding previously undescribed units where necessary. Forty-two community types were identified and mapped at Level 3. At Level 6, 90 different community types were mapped. The total area mapped was 35.2 km<sup>2</sup>, of which 31 km<sup>2</sup> were within the borders of BISC.

By providing a spatial inventory of the plant communities within the BISC, this map, along with existing data, will allow resource managers to effectively focus their restoration efforts and resources on communities that are indicative of relic or pristine conditions, or communities that are likely to benefit the most from active management. Furthermore, this map, with its 1:300 scale shoreline, serves as a turn-of-the-century baseline for the extent of mangroves within Biscayne National Park and, as a result, can be used to monitor the effects of sea-level rise on the wetlands and forested communities of Biscayne National Park for years to come.

## Introduction:

Established as a National Park in 1980, Biscayne National Park (BISC) is the largest marine park in North America. It comprises an area of nearly 700 km<sup>2</sup>, of which 95% is under water (Figure 1). The remaining 5% consists of uplands embedded within the Biscayne Bay Coastal Wetlands (BBCW), a network of freshwater and brackish wetlands along the western shore of Biscayne Bay, and within a set of Key Largo limestone (Stephenson & Stephenson 1950) barrier islands that parallel the mainland several kilometers offshore and define the eastern rim of Biscayne Bay (Figure 1). While these terrestrial communities are limited in area, about 31 km<sup>2</sup>, they extend over many kilometers and account for a significant portion of the Park's biodiversity, which includes many threatened and endangered species of both flora and fauna. Mangrove communities (forest, woodland, shrubland, and scrub) are by far the most abundant vegetation type found within BISC. Coastal Hardwood Hammocks are another important community type found within BISC, which are often overlooked. However, in contrast to the mangrove communities that are found throughout the Park, Coastal Hardwood Hammocks are restricted to the highest elevations within 11 of 42 mangrove-dominated islands found in BISC.

Not unlike other natural areas under the stewardship of the National Park System, BISC has endured many years of neglect and abuse by homesteaders and developers who, prior to the establishment of Biscayne National Monument in 1968, had free reign of the area and in many cases successfully drained, cleared, farmed, and developed much of these environmentally sensitive and ecologically important lands. During the latter part of the 1800's, pineapple and lime plantations were common throughout the Florida Keys including many of the islands now part of BISC (Leynes and Cullison 1998). Along with pineapples and limes, other fruits and vegetable were grown on the highly organic and productive soils associated with the Coastal Hardwood Hammock community. As a result, many acres of virgin Coastal Hardwood Hammock were cleared for cultivation and homesteads within BISC (Leynes and Cullison 1998). In the 1920's, Boca Chita, Adams, and Elliott Key were enlarged (Leynes and Cullison 1998) as developers sought to take advantage of the growth and prosperity of Miami Beach and the




Figure 1: Boundary of Biscayne National Park, Homestead, FL., USA.

surrounding areas as a vacation resort. In conjunction with the homesteaders and on the bequest of developers, public works projects dating back to the early 1900's for the purpose of mosquito control, land reclamation, and storm surge protection combined to compartmentalize and further alter the natural vegetation of BISC (Ruiz & Ross 2004).

Fortunately, 28 years of mitigation efforts have significantly improved the conditions of BISC. However, as a consequence of these historical land management practices, Biscayne Bay and the BBCW have become hydrologically isolated from the interior freshwater watershed that once flowed freely by sheet flow from the Everglades through the transverse glades (Ruiz & Ross 2004). As a result, there has been a marked decrease in the volume and kinetics of freshwater runoff into Biscayne Bay via tidal creeks and springs (Gaiser et al. 2005). This, in turn, has altered the natural seasonal variability of surface water salinities throughout the coastal wetland ecotone of BISC (Gaiser et al. 2005). All of these factors, in conjunction with the steady rise in sea level over the last century (2.2 mm/yr, Ross et al. 1994), have yielded large-scale changes in the composition and structure of the vegetation communities within the mainland portions of BISC. The offshore upland communities of BISC have benefited from their general isolation from the mainland and appear to have, for the most part, returned to a natural state. However, there is still clear evidence of historical anthropogenic perturbations throughout these islands.

Topo-lithographic maps of the Park based on 1928 aerial photographs date back to the early 1930's (Figure 2). Though not intended to be used as vegetation maps, these 1:20,000 scale topographic sheets are likely the first comprehensive vegetation maps that exist of BISC. These




Figure 2: 1931 topo-lithographic sheet of Old Rhodes Key and Totten Key vicinity based on 1928 aerial photographs.

topographic sheets depict the vegetation of the BBCW and BISC as a mosaic of broad scale community types characterized by deciduous hammocks, high ground deciduous (forests), mangroves, sawgrass on rock, and sloughs and document the extent of anthropogenic alteration in these natural areas for the period. Other maps of BISC include the 1943 and 1967 maps of Davis, the 1999 Florida statewide water management district land cover map, and the 2007 National Wetland Inventory map, among others. Unfortunately, all of these mapping efforts failed to capture the natural complexity and structure of the vegetation communities of the BBCW and BISC, in particular. To date, the only two mapping efforts which have succeeded at capturing the nature of the vegetation within BISC are the 1999 vegetation map of Welch et al. (1999) (Figure 3) and the 2002

vegetation map of Ross and Ruiz (2003) (Figure 4). Unfortunately, the Welch et al (1999) map is plagued with many commission and omission errors, is nearly 10 years old, and represents the conditions of BISC shortly after Hurricane Andrew (August 24, 1992). The 2002 Ross and Ruiz (2003) map, on the other hand, while spatially and thematically accurate, was limited in area and only included a small portion of the Park along its western shore (Figure 4). In 2006, because of the lack of a current, accurate, and useful vegetation map of BISC, the National Park Service Inventory and Monitoring Program South Florida / Caribbean Network commissioned the development of a spatially and thematically accurate vegetation map of BISC. This map, when completed, will provide a spatial inventory of the plant communities within the Park's jurisdiction with a level of accuracy suitable for planning, implementing, and quantifying management decisions and restoration efforts for the next several decades.




Figure 3: Welch et al. (1999) vegetation map of Old Rhodes, Totten, and Little Totten Keys, Biscayne National Park, Homestead, FL., USA.




Figure 4: Ross and Ruiz (2003) vegetation map of the Biscayne Bay Coastal Wetlands between the Princeton and Mowry Canals, Homestead, FL, USA

## Methods:

This project called for the creation of a spatially and thematically accurate vector map of the terrestrial communities of Biscayne National Park. As a result, a map was developed using traditional aerial photo-interpretation techniques, in conjunction with *Definiens Professional*® v5 remote sensing software (Definiens Imaging, 2006), NDVI calculated from the Red and NIR bands of each orthophoto, and LiDAR available for mainland portions of the Park through Florida International University (IHRC 2004). Vegetation communities or polygons were screen-digitized using ESRI® ArcMap™ 9.2 and stored in a Personal Geodatabase. The map is projected to NAD83, UTM Zone 17N (units = meters). The minimum mapping unit was set to 400 m<sup>2</sup>, but notable objects smaller than 400 m<sup>2</sup> were sometimes mapped based on the photo-interpreter's discretion. Digitized polygons were classified to the highest feasible level of resolution within the six-tiered hierarchical vegetation classification system developed by Rutchey et al. (2007) v5.22.07. Level 3 of the hierarchy was the minimum resolution required for this mapping project. However, some communities were mapped to Level 6. This map is expected to have a classification accuracy, at Level 3, of no less than 80% with 90% confidence. In addition, the expected minimum positional accuracy for well-defined objects was set at ± 5.0 meters of their actual location.

### *Imagery:*

The Fish and Wildlife Research Institute – Florida Fish and Wildlife Conservation Commission, provided the imagery used in this project. The imagery consists of a set of 316 images taken by Photo Science in May 2005. The images are a 5-band (RGBNIRPan) stacked multispectral orthophoto derived from three separate products. The first consist of a Red-Green-Blue (RGB) product; the second a Color-Infrared (RGNIR) product from which the redundant red and green bands were dropped; and the third a panchromatic (Pan) product. According to the metadata, the images were flown at an averages altitude of 3,142.2 m which produces an image scale of 1:26,185. Each image covers an area of about 10.5 km<sup>2</sup> with an average overlap of approximately 23% between adjacent images. The images are projected to NAD83 UTM Zone 17N and have a spatial resolution of 0.3048 meters. Regrettably, these images are neither spectrally calibrated nor geometrically correct. The calibration issue is the more serious problem in this project, since the lack of spectral correction results in glare and white-outs on bare-ground and water, and in identical vegetation communities having different spectral properties within and across images (Figure 5). The lack of geometric correction caused objects to lean outward from the principal point of the image, even though these images were visually checked against themselves, DOQQs, and different shorelines for planimetric accuracy. As a result, overlapping scenes or pixels from adjacent images were not coincident.



Figure 5: Spectral difference between overlapping adjacent images at 2 standard deviations.

### *Vegetation Classification System:*

Vegetation communities within BISC were classified to the highest feasible level of resolution within the six-tiered hierarchical vegetation classification system developed by Rutchey et al. (2007) v5.22.07. Level 3 (**L3 - Group**) of the hierarchy is the minimum resolution required for this mapping project. However, some communities were mapped to Level 6 (**L6 – Alliance**). Not surprisingly, not all communities registered on the orthophotos, observed in the field, and mapped were listed within this classification system. At the same time, many of the communities listed in the Rutchey et al. (2007) classification system were not observed within the boundaries of BISC. As a result, the classification system used in this project was modified to include all new community types observed and mapped and excluded types not present within the mapping area. Twenty-one new plant community types were added: 7 at Level 3, 5 at Level 4, 8 at Level 5, and 1 at Level 6 (see Appendix 1). Three other modifications were introduced to the Rutchey et al. (2007) classification system.

The first of these was the renaming and restructuring of the original Level 1 *Non-Vegetative* Class to *Other*. This new Level 1 *Other* class was then subdivided into two separate categories at Level 2: *Non-Vegetative* and *Anthropogenic*. This restructuring permits the discrimination, at Level 3, between non-vegetative natural environments (e.g. *Beach, Mud, & Water*) and environments, either vegetated or not, (e.g. *Lawns & Landscaping, Agriculture, Parking Lots, & Road*) that result from anthropogenic activities. The second modification was the standardization, wherever possible, of Raster ID's so that a species or community type would always be represented by the same number combination. For example, in the Rutchey et al. (2007) classification system, the following Raster ID's 114000, 315000, and 414000 represent the following three red mangrove communities': Forest, Shrubland, and Scrub, respectively.

By standardizing the 4<sup>th</sup> digit of each Raster ID, in this case to 4, we are able to discern quite easily that the following Raster ID's: 114000, 214000, 314000, and 414000 all represent a community type dominated by red mangrove. The third modification was the removal of blank categories between subordinate levels in the classification system. Aside from these exceptions, the new abridge version of the Rutchey et al. (2007) classification system (Appendix 1) keeps true to the original structure and hierarchy. However, since the Rutchey et al. (2007) classification system was design as an all-encompassing vegetation classification system for Southern Florida it was necessary, in some cases, to append the original community descriptions so that they would be less ambiguous and more relevant to this project (see descriptions below).

At the highest level of the hierarchy, Level 1 (**L1 – Class**), the new abridged classification system (Appendix 1) has the following structure and requirements:

- **Forest:** High-density stands of trees (>50% tree canopy cover) with heights > 5 meters
- **Woodland:** Low-density stands of trees (10 - 60% tree canopy cover) with heights > 5 meters in a matrix of shrubs, graminoids, and/or herbaceous vegetation.
- **Shrubland:** Stands of small trees and/or shrubs (canopy cover  $\geq$  50%) with heights < 5 meter tall.
- **Scrub:** Communities of dwarf trees or shrubs typically in a matrix of graminoids, and/or herbaceous vegetation. Canopy cover 10% to 50% but can be as high as 100% for Mangrove. Canopy < 5 meters tall with the exception being for Mangrove which is  $\leq$  2 meters.
- **Marsh:** Graminoid and/or herbaceous emergent or floating vegetation in shallow water that stands at or above the ground surface for much of the year.
- **Dune:** A ridge of wind blown or windstorm deposited sand or similar material directly inland and parallel to the shoreline which is commonly vegetated by graminoids and/or herbs and sometimes even shrubs.
- **Submerged Aquatic Vegetation:** Vegetation that has evolved the ability to carry out its entire life cycle completely submerged in an aquatic environment.
- **Exotic:** Non-native and invasive vegetation.
- **Other:** Non-vegetative or anthropogenic cover.

The next level in the hierarchy, Level 2 (**L2 – Type**), modifies the structurally-defined Level 1 with a community designation, for example, *Mangrove Forest* vs *Hammock Forest* or *Mangrove Shrubland* vs *Upland Shrubland*. At Level 3 (**L3 – Group**) the classification system often requires that the previous level (**L2 – Type**) community types be identified by dominant species, such as *Black Mangrove* or *Buttonwood Forest* or *Red Mangrove* or *White*

*Mangrove Shrubland*, for example. However, this is not always the case. For example, at Level 3, *Hammock Forest* can be subdivided into *Coastal Hardwood Hammocks* or *Coastal Dune Hammocks*. A similar situation occurs within the Level 2 *Salt Marsh* and *Freshwater Marsh* types. At Level 3, these two types are differentiated based on morphological traits like graminoids, herbaceous, or succulent for Salt Marsh or by Marsh or Prairie for Freshwater Marsh. The remaining three levels of the classification system (**L4 - Formation**, **L5 – Alliance**, & **L 6 – Association**) continue to subdivide the previous community types by dominant canopy species composition and then by understory species assemblages; for example, a Level 6 mangrove scrub community might have the following nomenclature *Buttonwood-White Mangrove Scrub-Glasswort*.

*Shoreline Segmentation – Definiens Professional® v5:*

*Definiens Professional® v5* remote sensing software was incorporated into this project as an exploratory methodology for standardizing and automating the digitization of vegetation communities into polygons. For mapping vegetation, *Definiens Professional® v5* was mostly unproven but offered tremendous benefits as a remote sensing tool. In contrast to traditional remote sensing software in which an image is classified at the pixel level, *Definiens Professional® v5* classifies images in terms of image objects and their mutual relationships. In other words, pixels with similar brightness values and/or other characteristics are aggregated into image objects. These image objects are then the target of the classification process and not the pixels themselves. Image objects within *Definiens Professional® v5* are derived by one of three distinct segmentation algorithms: chessboard segmentation; quadtree based segmentation; or multiresolution segmentation. For this project, all image objects were derived using the multiresolution segmentation. This algorithm is a heuristic optimization procedure that locally minimizes the average heterogeneity of image objects for a given resolution (Definiens 2006). Image objects derived from this methodology are created based on three criteria: scale, color, and shape. The scale parameter is an abstract terms that determines the maximum allowed heterogeneity within image objects and their average size (Definiens 2006). Color and shape are mutually exclusive weighted parameters that determine which heterogeneity attributes are minimized during the segmentation process (Definiens 2006). Color emphasizes how brightness values within layers will contribute to the entire homogeneity criterion. On the other hand, shape, composed of two weighted parameters: smoothness and compactness, defines the textural homogeneity of the image objects by either optimizing for smoothness or compactness. Segmentations weighted exclusively towards color result in spectrally homogeneous but spatially heterogeneous image objects, while, segmentations weighted towards shape result in image objects that are spatially homogeneous but spectrally heterogeneous. Regardless of the color weighting, image objects tend to follow natural features more closely when smoothness is emphasized over compactness (Benz et al. 2004).

Multiple exhaustive attempts where made using *Definiens Professional® v5* to produce image objects that could be classified into distinct vegetation communities. However, it became clear, early on, that these segmentations efforts were accentuating the spectral variability of the imagery and not yielding objects that could be classified into distinctive vegetation

communities based on a homogeneous spectral signature. Our efforts, however, showed that the land-water interface could be effectively and consistently segmented into image objects to create fine scale (1:300) shorelines.

Two similar, yet contrasting, methodologies were developed for the delineation of the land-water interface. The first methodology (Appendix 2) is a multifaceted segmentation algorithm modeled from the initial segmentation efforts designed to classify the varied vegetation communities within BISC. This methodology is described in Appendix 2. The second methodology was developed later in the project to: 1) quickly and efficiently recreate shorelines where the original segmentation was based on an image that was no longer deemed appropriate because of planimetric errors or unsuitable color balance; and 2) capture and preserve the fine scale details associated with small irregular mangrove islands or individual trees found on intertidal zones or shallow lagoons. This methodology, described below, while simplistic and straight forward, was as robust as the original segmentation algorithm and minimized the number of image objects needed to consistently segment land and water into image objects.

To minimize file size and processing time, only the Red and NIR spectral bands of each image were imported into *Definiens Professional*® v5. The NIR band was used to define the no data area. Based on the brightness values of the imported two bands the ratio NDVI (see NDVI methods section for equation and description and used of this ratio) was defined as an arithmetic custom feature. This ratio was later used in the sample editor to help classify the image objects. The next step was to define the scale, color, and shape parameters of the multiresolution segmentation. Because of the lack of spectral calibration, the scale parameter for each image segmentation was determined empirically. In general, the larger the scale parameter used the larger the image objects created. At the same time, the larger the scale parameter the fewer image objects created and the less time it takes to run a segmentation. As a result, the initial scale parameter for all segmentations were set to 100. Subsequent segmentations, if needed, were decreased systematically until they reached a minimum value of 10. Scale parameter values < 10 were impractical with this imagery because of the amount of processing time needed to perform the segmentation and the overwhelming number of image objects created, about 260,000 at this scale. In all cases, the Color parameter was set to 0.8, forcing the Shape parameter to 0.2, and the Compactness and Smoothness were set at 0.2 and 0.8, respectively. After the multiresolution segmentation was completed, a spectral difference segmentation was run to merge spectrally similar adjacent objects. In general, values between 2 and 5 were used. Values greater than 5 usually reduce the number of image objects by half but, as a result, merged too many image objects together and thus created many mixed image objects unsuitable for classification. The next step was to classify the image objects. Classification of the image objects was defined through a class hierarchy dialogue similar to Method 1 (Appendix 2). However, only two classes were used: Water and Land. The Water class has an explicit definition (NDVI less than an optimal value determined empirically for each image), whereas the Land class was defined as those image objects not classified as Water. Once the image objects were classified and the classification validated, the image objects were converted to polygons and merged into their respective classes; Water and Land. These merge features were then exported as a vector shapefile for final editing.

Within ESRI® ArcMap™ 9.2 the newly imported shapefile was converted into a grid and then back-converted into a shapefile for final validation and editing at a scale of 1:300. The reasoning behind this double transformation is as follows: Segmented image objects in *Definiens Professional*® v5 tend to be highly fractal (Figure 6a). The transformation to a grid and then back to a polygon has the effect of removing unnecessary vertices from the polygons and thus smoothing the overall shape of the polygons into slightly more realistic and aesthetically pleasing polygons (Figure 6b). Another advantage of this smoothing transformation is that by smoothing the overall shape of the polygons, shoreline errors that would normally have had to be manually corrected are, for the most part, processed automatically, thus saving precious time in the final editing session.




Figure 6: Contrast between shorelines without (A) and with (B) the smoothing transformation. See methods section for methodology.

#### *NDVI:*

The Normalized Difference Vegetation Index (NDVI) ratio was calculated for each image as a 1-band gray-scale raster in Erdas Imagine 9.1. NDVI is calculated by subtracting the Red spectral band from the NIR spectral band and then dividing this difference by the sum of the NIR and Red spectral bands (Equation 1).

$$\text{Equation 1: } \text{NDVI} = (\text{NIR} - \text{Red}) / (\text{NIR} + \text{Red})$$

Technically, NDVI should be calculated on the corrected radiant flux of each pixel. However, since none of the imagery was spectrally calibrated NDVI was instead calculated using the original brightness values of each pixel. As a result, community trends in NDVI values could only be applied to the imagery it was derived from and not across all images uniformly. In

general, however, NDVI values  $> 0.3$  usually corresponded to heavily vegetated areas with dense canopies characteristic of Shrubland and Forest communities (Figure 7). Scrub communities were usually associated with NDVI values between 0.1 and 0.3 (Figure 7). Soils and bare ground, have a higher reflectance in the near-infrared than in the red and thus exhibited low NDVI values generally between  $-0.1 - 0.1$  (Figure 7). Water, on the other hand, because of its low reflectance in both spectral bands (Red and NIR), tended to have very low NDVI values, usually  $< -0.1$  (Figure 7). Base on these criteria, NDVI helped delineate low productivity zones (Mangrove Scrub) and Non-Vegetative features from adjacent Shrubland and Forest communities. However, NDVI failed to discriminate between the more productive community types (e.g. Hardwood Hammocks, Mangrove Shrublands, and Forest).




Figure 7: General relationship between community type and NDVI values.

### *LiDAR:*

Light Detection and Ranging (LiDAR) is an active airborne remote sensing technology that uses laser pulses to provide direct and accurate measurements of vegetation structure and topography, among other things. Modern airborne laser systems have demonstrated the ability to: quantify structural changes in mangrove forests following hurricanes (Zhang et al. 2008); provide accurate estimates of forest vertical structure and volume (Zhang et al. 2006, Suárez et

al. 2005), and stem diameters (Hyypä et al. 2001); and have outperformed other remote sensing systems at estimating mean tree height (Lefsky et al. 2001). In recent years, LiDAR has become an invaluable remote sensing tool with many practical applications including land cover classification and mapping (Bork & Su 2007, Hill & Thomson 2005).

The LiDAR used in this project was provided and flown by Florida International University International Hurricane Research Center (IHRC 2004). This data was part of a 2002-2003 survey of Miami-Dade County, Florida. However, the extent of the LiDAR coverage for BISC was, unfortunately, limited to the uplands on the western shore of Biscayne Bay starting just north of Turkey Point and did not include any of the islands within BISC. The flight and laser acquisition parameters (altitude, speed, and scan angle, scan rate, and pulse rate) produced, on average, a 650-meter-wide swath of 30 cm wide laser footprints with a nominal point spacing of 1.5 m (IHRC 2004). The first return (first-stop) data from this survey was interpolated using kriging to produce 5 ft (1.524 meters) resolution top-surface digital elevation models (DEMs) with a vertical accuracy of  $\pm 0.80$  ft (24 cm) (IHRC 2004). For BISC, these DEMs represent the mean canopy height (ft) of the vegetation communities sampled in 2002 (Figure 8). The DEMs are projected to State Plane, Florida East Zone, NAD83 (unit = feet) with NAVD88 (feet) as the vertical datum (IHRC 2004).




Figure 8: 2002 top-surface digital elevation model (DEM) of LiDAR data for Convoy Point, Homestead, FL.

Since the LiDAR was collected in 2002, the DEMs had to be corrected for canopy height growth between 2002 and 2008 (Table 1). Changes in canopy height were scaled according to community type. Thus, extrapolation of canopy heights was weighted towards the more productive forest community where tree height growth rate can easily reach and exceed 1 m/yr. In contrast, the less productive shrubland and scrub communities would not be expected to have canopy height growth much higher than 20 cm/yr or 5 cm/yr, respectively (Ross et al. unpublished data). The canopy height correction to the DEMs was done in ESRI® ArcMap™ 9.2 as a custom natural breaks classification. Thus, the original DEM values (ft) were reclassified and labeled to show the predicted canopy height in meters for 2008. The classification was saved as a layer file and imported as a symbology each time a new DEM was used.

Table 1: Extrapolation of canopy LiDAR height (m) between 2002 and 2008 used in discriminating between Level 1 community types.

<b>Community Type</b>	<b>2002 LiDAR Height (meters)</b>	<b>2008 Extrapolated LiDAR Height (meters)</b>
Non-Vegetative	< 0	< 0
Scrub	0 – 1.75	0 – 2
	1.75 – 2.1	2 – 2.7
Shrubland	2.1 – 3	2.7 – 3.7
	3 – 3.9	3.7 – 4.9
Forest	> 3.9	> 4.9

#### *Unsupervised Classification:*

In an exploratory manner, two independent unsupervised classification methods, pixel based and image-object based, were used in this study to ascertain their potential to classify the vegetation communities of BISC. The area used in this analysis had been previously mapped by Ross and Ruiz (2003) and thus could be use to quantify how well these two methods worked at delineating community boundaries.

In all, three 11-band stacked images were used in this analysis. The 11-band stacked images were derivatives of the original set of images provided for this study and were created using Erdas Imagine 9.1. The first five bands included the Red, Green, Blue, NIR, & Panchromatic bands from the original images, resampled to 0.9 meters. The 6<sup>th</sup> band was a 3 meter resolution gray-scale NDVI layer. The remaining 5-bands, #'s 7-11, consisted of the five bands from the original images, resampled to 3 meters using a 3x3 fixed window variance filter. Images were cropped to remove areas that contributed excessive glare or were not relevant to the study area. This eliminated unnecessary data from the analysis that might interfere with the classification.

The pixel-based unsupervised classification was carried out using the ISODATA algorithm in Erdas Imagine 9.1. The 11-band image stack described above was resampled to 9 m for this classification, however. A visual inspection of several image resolutions suggested that this scale of resampling preserved the details of the vegetation landscape, while, reducing the ‘scatter shot’ appearance of individual single pixels of many classes somewhat. The ISODATA algorithm was applied to all 11 bands to produce 40 classes. Subsequently the classification

was opened in ESRI® ArcMap™ 9.2, and classes were combined to produce homogeneous areas that best matched the original resolution of the images (0.3048 m) or the Ross and Ruiz (2003) vegetation map.

The image-object unsupervised classification was initiated in *Definiens Professional*® v5 using the same 11-band image stack described above. A multiresolution segmentation was run using 9 of the 11-bands in the image stack and on a shoreline vector layer, previously created in *Definiens Professional*® v5. While adjusting for scale, several segmentation runs were done. Spectral information was weighted over shape by using a shape factor of 0.2. To reduce the appearance of irregular jagged objects, the smoothness parameter was weighted at 0.8 compared to 0.2 for compactness. Several complete analyses indicated that the size of the image objects created in *Definiens Professional*® v5 has a large impact on the quality of the final product. If objects are too small, they only reflect local variation and not the pattern of the vegetation community. The best classification results were seen with approximately 1,800 image-objects using a scale parameter of 200. This segmentation was then exported to a vector polygon file consisting of 30 features representing spectral information, size, shape, and texture.

Many of the exported features were highly correlated. To improve the performance of the subsequent principal components analysis (PCA), features were removed so that correlations between them did not typically exceed 60 percent. Representative features used in the PCA included: area, roundness, compactness, shape index, mean object NDVI pixels in object, mean of variance filtered NIR pixels, mean of variance filtered panchromatic pixels, mean of red pixels in object, mean of green pixels, mean of blue pixels, mean of NIR pixels, mean of panchromatic pixels, SD of NDVI pixels in object, ratio of object length to width, max difference, density and asymmetry.

PCA on the attributes of the exported image objects was carried out using Proc PRINCOMP in SAS v9. Based on previous results, the first 4-5 axes of the PCA usually represent about 80-85% of the total variability. As a result, these axes were used in the subsequent cluster analysis. A cluster analysis was also carried out in SAS using the K-means clustering algorithm, Proc FASTCLUS. Many of the 20-40 classes found represented non-vegetation areas, such as artifacts in the boundaries of the imagery and water areas. In one case, seven of 30 classes representing mixed vegetation were reclassified into 20 additional classes. Once the PCA and cluster analyses were complete, the attribute table was merged back into the object geometry shapefile in ESRI® ArcMap™ v9.2. A visual assessment of the unsupervised classification was done by comparing object boundaries and classes with the Ross and Ruiz (2003) vegetation map and with the original aerial photographs of the study area.

#### *Geodatabase Design & Mapping:*

Currently, geodatabases are the most efficient way of editing and storing spatial information. Geodatabases merge traditional GIS data formats and data management tools to create a geospatial environment, which maintains consistency and accuracy by defining how data is stored, accessed, modified, and managed within a single file. For this project, a Personal

Geodatabase, containing 19 feature datasets and 5 feature classes, was created in ESRI® ArcCatalog™ v9.2. All feature datasets and classes within the Geodatabase are projected to NAD83, UTM Zone 17N (unit = meters).

Each feature dataset within the Geodatabase represents an island or region within BISC and contains a topology and two feature classes, *shoreline* and *vegetation*. The naming convention for each topology and feature class is defined by using the feature dataset name as the prefix and the topology and feature class type, *shoreline* or *vegetation*, as the suffix (e.g. , *Adams\_Key\_Topology*, *Adams\_Key\_Shoreline* or *Adams\_Key\_Vegetation*). Within each feature dataset, the topology contains three rules that ensure the proper management of coincident geometry between polygons and the *shoreline* and *vegetation* feature classes. The *shoreline* feature class defines the land-water interface. The structure of this feature class remains consistent throughout all feature datasets and contains the following default ESRI attribute: OBJECTID, SHAPE, SHAPE\_Length (m), & SHAPE\_Area (m<sup>2</sup>). The structure of the *vegetation* feature class is also constant throughout the geodatabase and contains the following attributes for each record: OBJECTID, SHAPE, AUTHOR, QC, RASTER ID, L1, L2, L3, L4, L5, L6, Photo, SHAPE\_Length, & SHAPE\_Area (Table 2).

Table 2: Description of attributes for vegetation feature class.

Attributes	Description
OBJECTID	ESRI default (ID)
SHAPE	ESRI default (polygon)
AUTHOR	Person responsible for the creation or last edit of that record
QC	Date used to designate when the record was verified in the field or to suggest a high degree of confidence regarding its classification on the day it was digitized
RASTER ID	Numeric string representing the mapped vegetation at the lowest level of the classification
L1	Level 1 classification
L2	Level 2 classification
L3	Level 3 classification
L4	Level 4 classification
L5	Level 5 classification
L6	Level 6 classification
Photo	Orthophoto used to create this record
SHAPE_Length	ESRI default (shape perimeter m)
SHAPE_Area	ESRI default (shape area m <sup>2</sup> )

The five, additional, independent feature classes in the geodatabase; i.e. *BISC\_Boundary*, *GR\_Points*, *OrthoPhoto\_Index*, *Shoreline*, & *Vegetation*, represent miscellaneous thematic layers that define the extent of the map, document the data collected, and summarize the spatial information populated within the geodatabase. The *BISC\_Boundary* feature class contains one record denoting the 2006 legislative boundary of Biscayne National Park. This feature was acquired directly from The National Park Service Inventory and Monitoring Program South Florida / Caribbean Network. *GR\_Points* is a point feature class populated with the 1,081 ground-reference data points collected to assist in the identification of unknown types or regions within the imagery or to ground-truth communities and areas already mapped. This feature class contains both structural and species information. *OrthoPhoto\_Index* is a

polygon feature class that serves as the photo index for this project. It documents which orthophoto was used to delineate the shoreline and map the vegetation for a particular region. Knowing and documenting this information was crucial because the imagery provided were not geometrically correct, as discussed previously. Consequently, digitized features not paired with the originating orthophoto could have, on average, a spatial offset of 2 meters or more. The last two feature classes, *Shoreline* and *Vegetation* contain all of the data within each of the 19 feature datasets and are the master Shoreline and Vegetation feature classes.

Edits to all feature classes were done within the geodatabase environment in ESRI® ArcMap™ v 9.2. The *Definiens Professional*® v5 shorelines were edited using the 2005 original imagery at a scale of 1:300. The vegetation communities, on the other hand, were delineated (digitized) at a 1:1500 scale using the 2005 imagery. Delineation of the vegetation boundaries was facilitated by overlaying the imagery, with a 25% transparency, over the NDVI raster or the LiDAR DEMs (Figure 9A & 9B, respectively).




Figure 9: NDVI (A) and LiDAR (B) overlapped by imagery with 25% transparency.

#### *Accuracy Assessment & Ground-truthing:*

The accuracy assessment of this map will be subcontracted to an independent contractor within 12 months of project completion. This will ensure that the integrity of the map is not compromised and that the minimum standards set forth by the cooperative agreement between the National Park Service and Florida International University, for this project, are met.

Over 1,000 ground-reference points accessed by helicopter, boat, and on foot (Figure 10) were used to ensure that the map would meet the expected accuracy of 80% with an allowable error of 10% at Level 3 of the classification. On average, 120 individual sites were visited per day. A total of 9 field days were used on this endeavor; four of which were helicopter days. Field observations included, but were not limited to, maximum canopy height, canopy species composition, herbaceous layer species composition, and, on occasion, substrate. The GPS locations of each site visited was recorded directly into a field-book using a handheld GPS or populated into a geodatabase directly via a submeter accurate GIS/GPS integrated system. This information was then used to interpret the spectral signature of unknown types based on the spectral signature of known communities verified by the ground-referencing locations.




Figure 10: Distribution of ground-reference points used to map Biscayne National Park.

## Results & Discussion:

### *Vegetation Classification System:*

One of the unstated goals of this project was to determine the potential and effectiveness of the Rutchey classification system at mapping and capturing the natural complexity and structure of the varied plant communities arrayed along local environmental gradients in BISC. To that end, we found the classification system adequate. However, inherent issues with the classification system, particularly within the mangrove subcategories, created issues that necessarily lead to commission & omission errors. Providing a sound solution to these issues is beyond the scope of this project and report. Nonetheless, the following are worth noting:

- The classification system relies heavily on knowing the following three parameters: height, density, & species composition. Height and density are the most important at the highest level of the hierarchy; i.e. Level 1. Conversely, species composition becomes important at the 3rd level of the classification and thereafter. Height and density are usually difficult and at times impossible to discern from most aerial photography unless one of the following criteria are met: 1) the imagery is of high enough quality and has a spectral resolution that allows for individual crown to be identified; 2) the imagery consists of stereoscopic pairs and a stereoscope is used; or 3) there is enough textural contrast and shadow between communities that their vertical structure can be inferred. As a result, if the imagery, along with the methodology used, fails to recognize and meet the criteria described above, a situation is created where classification errors can occur at a level within the classification system (i.e. Level 1) where they should not be occurring. These errors are problematic since they are then inherited by the subordinate categories. Few can argue that there is a clear solution to this problem. However, the incorporation of LiDAR can mitigate and significantly, if not completely, eliminate classification errors within the 1<sup>st</sup> Level of this classification system and thus increase the overall accuracy of a map.
- Based on height and density, the Exotic Level 1 Class should be merged into the corresponding 1<sup>st</sup> Level structural class (i.e. Forest, Woodland, Shrubland, or Scrub). For example, a greater than 5 meter tall stand of Australian Pine with greater than 50% cover should be assigned to the L1 – Class: Forest with the following or similar hierarchical structure:

L1: Forest

L2: Non-Native Forest (or similar description)

L3: Australian Pine (*Casuarina* sp.) Forest

L4: subcategory if applicable (specific species)

L5: subcategory if applicable (treated or not treated)

L6: subcategory if applicable

This reorganization is logically consistent to the overall organization of the Rutchey et al. classification system and is applicable to all non-native community types. Moreover, this

hierarchical structure emphasizes the nature and structure of the non-native community. By documenting these communities based on their canopy structure, resource managers are better able to properly ascertain the amount of resources needed for exotic control teams to effectively combat, eradicate, and mitigate these communities.

Co-dominance between native and non-native species and the multi-structural (stratum) characteristics of plant communities are related issues that should be addressed, in general as well as in BISC. For example, it is not uncommon for Australian Pine to be associated with a mixed mangrove community (e.g. Buttonwoods and White mangroves) in a Forest setting in which neither association (i.e. mangroves or non-natives) is dominant. Other examples include dense mixed-mangrove shrublands overtopped by a sparse or full canopy (30-75% cover) of Australian Pine, which are commonly associated with both mosquito & drainage ditches; Australian Pine shrubs or individual trees in a matrix of sawgrass (*Cladium jamaicense*); or mangrove forests (mixed or monotypic) with a dense understory of Brazilian Pepper (*Schinus terebinthifolius*) or Shoebutton Ardisia (*Ardisia elliptica*). These examples may seem like extreme cases rarely seen, but in actuality are commonplace throughout many of the interior wetlands located on the western shore of Biscayne Bay. Unfortunately, deriving a classification which can properly address these complexities is not easy and requires a significant amount of further thought and discussion by the authors and end users of the Rutchey classification system.

- At the 3rd level of the mangrove hierarchy, regardless of class (i.e. Forest, Woodland, Shrubland, Scrub), the cartographer or photo-interpreter is faced with the challenge of identifying mangrove communities at species level (e.g. Black Mangrove Forest, Mixed Mangrove Shrubland, or Red Mangrove Scrub, etc.). In many instances, it is not an easy task to distinguish dominant species (e.g., Black Mangrove, Red Mangrove, Mixed Mangrove), and commission and omission errors are commonplace. In regard to the overall organization of the other community types within the classification system (e.g. the Marsh category, which contains a complete hierarchical structure), it would seem that this species level of detail is better suited for the next lower level (**L4 - Alliance**) or lower within the classification system. The reason for this is two fold. For one, it would eliminate the gap in the mangrove hierarchy that currently exists between Level 3 and Level 5; i.e. there is no Level 4 to transition between many Level 3 (e.g. Red Mangrove Scrub) and Level 5 communities (e.g. Red Mangrove Scrub-Graminoid). The other reason is in the difficulty in determining the mangrove species composition from imagery alone at Level 3, particularly in the Shrubland and Scrub classes. Furthermore, there are many instances when the spectral signature of a lower level community associate like, Saltwort (*Batis maritima*), is readily recognized but the mangrove community associated with it is not (Figure 11). For example, in Figure 11, the succulent understory (yellowish in color) is clearly visible, while the species composition of the overtopping community, if one is present at all, is more difficult, if not impossible to distinguish. Thus, without auxiliary data this community has a high probability of being misclassified based on the overwhelming signature of Saltwort, thereby affecting the thematic accuracy of the map.




Figure 11: The succulent understory of the community outlined in blue (definable at Level 5 of the classification) is clearly visible, but the structure and species composition of the canopy, defined at Level 3, is difficult to determine. Based on a ground survey, this community was determined to be a Mixed Mangrove (Buttonwood-White Mangrove) Scrub with an understory dominated by Saltwort.

The solution to the two issues discussed in this bullet might be solved with the following modification to the classification system. The example below applies to Mangrove Scrub, but could apply equally well to other Scrub, Shrub, or Woodland categories:

**L2: Mangrove Scrub**

**L3: Mangrove Scrub-Closed Canopy**

L4: Black Mangrove Scrub

L4: Mixed Mangrove Scrub

L5: Black Mangrove-Red Mangrove Scrub

**L3: Mangrove Scrub-Succulent**

L4: Mangrove Scrub-Glasswort

L5: *White Mangrove Scrub-Glasswort*

L4: Mangrove Scrub-Saltwort

L5: *Black Mangrove Scrub-Saltwort*

L4: Black Mangrove Scrub-Succulent

L5: *Black Mangrove Scrub-Saltwort*

L4: White Mangrove Scrub-Succulent

L5: *White Mangrove Scrub-Glasswort*

L4: Mangrove Scrub-Mixed Succulent

L4: Mixed Mangrove Scrub-Succulent

L5: Buttonwood-White Mangrove-Succulent

L6: *Buttonwood-White Mangrove-Saltwort*

L5: Mixed Mangrove Scrub-Saltwort

L6: *Buttonwood-White Mangrove-Saltwort*

L4: Mixed Mangrove Scrub-Mixed Succulent

**L3: Mangrove Scrub Graminoid**

While the entire hierarchy is not displayed, the general structure and trend can be followed. By applying this hierarchical structure, the gaps between Level 3 and Level 5 are now gone. Moreover, from a remote sensing perspective, communities or individual plants or groups of plants that are readily discerned but buried deep in the classification hierarchy are now moved up into an appropriate level while communities with a complex and/or ambiguous spectral signature are demoted or grouped logically to minimize commission and omission errors. This restructuring of the classification system provides multiple pathways, at the lower levels of the hierarchy, for specific detailed community types to be arrived at while minimizing community misclassification at higher levels. Multiple pathways might appear problematic, but if the goal is to minimize classification errors and ensure a thematically correct map, while avoiding gaps within subcategories, this may prove to be the best solution within the parameters of the Rutchey et al. (2007) classification system.

#### *NDVI & LiDAR:*

As discussed in the methods section, NDVI was instrumental at delineating Non-Vegetative communities and low productivity zones from adjacent more productive communities. However, since the imagery was not spectrally calibrated, the interpretation of NDVI values shifted with each new image used. As a result, the use of NDVI was somewhat impractical because established rule sets had to be confirmed and calibrated each time a new image was used. Nevertheless, the potential of NDVI to distinguish communities along a productivity gradient was demonstrated in this mapping project.

In contrast, by providing direct and accurate seamless estimates of canopy heights for an area, LiDAR overcame the limitations inherent to NDVI and proved superior at distinguishing communities and their boundaries regardless of class and productivity. Consequently, the availability of LiDAR for the mainland portions of the park significantly improved the spatial and thematic accuracy for this section of the map. Moreover, LiDAR notably reduced the amount of time needed to interpret images and digitize plant communities. Furthermore, the multi-dimensionality of LiDAR (i.e. not only does it provide information regarding the mean canopy height for a vegetative community it also possesses information regarding vertical structure, stand volume, basal area, and on topographic relief, among others things.) contributes and enhance our understanding of the spatial and temporal dynamics of plant communities. As such, within the context of resource management, protection, and restoration LiDAR is an invaluable tool, which permits for the quick collection, analysis, and interpretation of large volumes of both spatial and temporal data over large areas without the logistical cost and difficulty associated with ground surveys.

*Unsupervised Classification:*

Figure 12 shows the pixel-based unsupervised classification using the ISODATA algorithm overlaid with the Ross & Ruiz (2003) vegetation map of the area, for comparison. In general, this classification outlines many of the community boundaries identified by Ross & Ruiz (2003). However, it failed to produce homogeneous areas characterized by one or two pixel classes, which are essential to properly isolate and delineate community types, particularly at the scale of this map. Since many of these classes appear closely related in space, the original 40 classes were consolidated into 10 new cover classes, which are described in Table 3. Taken as a whole, this new classification (Figure 13) matches more closely with the community boundaries drawn by Ross & Ruiz (2003). Nevertheless, the same ‘scatter shot’ effect which plagued the original classification remains. As a result, most communities still appear as a multi-pixel class aggregate making community delineation very difficult and impractical for this project.




Figure 12: Comparison of unsupervised classification with 40 classes overlaid with the Ruiz & Ross (2004) vegetation boundaries in black.



Figure 13: Comparison of unsupervised classification with 10 classes overlaid with the Ruiz & Ross (2004) vegetation boundaries in black.

Table 3: Description of unsupervised classification cover classes shown on Figure 12.

Class	Color	Description
1	Bright Blue	Water
2	Light Blue	Salt marsh and heavier scrub mangrove in areas between ditches.
3	Yellow	Scrub mangrove in areas of less vegetation west of class shown in light blue.
4	Green	Transitions to denser more textured vegetation.
5	Aqua	Coastal mangrove and some interior non-mangrove areas. This class is picking up areas of rougher texture for coastal and interior areas and when present in the fringe mangrove area is associated with the taller tree canopy.
6	Brown	Associated with rough textured areas near drainage ditches and the border with green and purple areas.
7	Purple	Associated with interior and transitional mangrove areas.
8	Black	Associated with drainage ditches, edges of water bodies, and tree shadows.
9	Red	Main association with <i>C. equisetifolia</i> forest in western part of image and coastal bordering fringe mangrove.
10	Orange	Lesser association with <i>C. equisetifolia</i> forest.

Image objects created in *Definiens Professional*® v5 containing a 17 factor attribute table were analyzed in SAS v9 using a principal components analysis (PCA) and K-means cluster analysis. The first five axes represented 84% of the variation observed in the data (Table 4). The 17 factors forming the basis of the PCA grouped into four distinct categories based on spectral and textural properties, shape, and a miscellaneous “other” category (Table 5).

Table 4: Eigenvalues for the correlation matrix

PC	Eigenvalue	Difference	Proportion	Cumulative
<b>1</b>	6.195	2.951	0.364	0.364
<b>2</b>	3.244	0.909	0.191	0.555
<b>3</b>	2.335	0.894	0.137	0.693
<b>4</b>	1.441	0.374	0.085	0.777
<b>5</b>	1.067	0.300	0.063	0.840
<b>6</b>	0.768	0.244	0.045	0.885
<b>7</b>	0.524	0.041	0.031	0.916
<b>8</b>	0.483	0.165	0.028	0.945
<b>9</b>	0.317	0.099	0.019	0.963

Based on the eigenvectors (Table 5), PC1 (36%) is a composite of several spectral measures. PC2 (19%) has a negative association with shape (Table 5). PC3 (14%) also appears to have a strong association with the overall shape of each feature. PC4 (9%) and PC5 (6%) are minor components.

Table 5: Eigenvectors (factor scores) computed for the correlation matrix.

Group	Factor	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5
Spectral	Red	0.286	0.273	-0.187	0.163	0.049
	Green	0.320	0.218	-0.238	0.205	0.002
	Blue	0.235	0.280	-0.251	0.332	0.025
	NIR	0.360	-0.121	-0.119	-0.154	-0.108
	Pan	0.365	0.124	-0.203	0.083	-0.020
	NDVI	0.280	-0.259	-0.093	-0.276	-0.140
Texture	SD NIR	0.151	-0.002	0.196	-0.373	0.500
	VarNIR	0.240	0.228	0.226	-0.296	0.164
	VarPan	0.252	0.266	0.087	-0.217	0.205
Shape	Area	0.146	-0.364	-0.023	0.162	-0.283
	Roundness	0.248	-0.308	0.221	0.225	0.236
	Compactness	0.144	-0.233	0.311	0.426	0.334
	Shape Index	0.252	-0.130	0.424	0.176	-0.113
	Ratio L to W	0.122	0.234	0.341	-0.175	-0.488
	Density	0.083	-0.328	-0.360	-0.314	0.177
	Asymmetry	0.133	0.118	0.314	-0.085	-0.249
Other	Max Diff	-0.263	0.324	0.116	0.124	0.233

In total, 1871 objects were classified into 30 classes using the first five principal components (Table 4) with K-means clustering implemented in Proc FASTCLUS. Seven of the original 30 classes (Class 6, 16, 17, 18, 19, 21 and 27) were found to be a mix of vegetation types that did not separate well into any one class. To improve the separability of these classes, a new cluster analysis was executed on only those objects (712) that grouped into these seven classes.

The results of this second cluster analysis are shown in Figure 14. Out of the 20 new classes, three had to be omitted because they were image artifacts. The results obtained from the remaining 17 classes reveal that the image object boundaries were not following the vegetation patterns visible in the aerial photographs, despite their small size and the inclusion of additional layers to capture texture information (variance filter) and to strengthen the signal (NDVI) between poorly and highly vegetated areas. Possibly, the scale parameter for the multiresolution segmentation was set too large. However, decreasing the scale setting would have created smaller objects that tend to be more jagged and abstract in shape, are heavily influenced by light and shadow, and thus are not representative of the community. In the




Figure 14: Comparison of object-based unsupervised classification with 17 classes overlaid with the Ruiz & Ross (2004) vegetation boundaries in black.

end, this unsupervised classification method also failed to properly classify and delineate community types within BISC.

In general, computer-based classification methods (pixel-based or object-based) of vegetation require that: 1) the imagery is radiometrically and geometrically corrected and 2) the spectral signature of the vegetation being classified is distinct enough to allow for its classification or separation into separate unique classes or vegetation communities, in this case. If these two parameters are not met, the computer-based classification will tend to have a high degree of commission and omission errors. In the case of this project, the imagery did not meet these standards and the vegetation types being classified did not have exclusive spectral signatures at Level 3 of the classification, the minimum required for this project. At the same time, the classification system applied (Appendix 1) relies heavily on knowing the canopy height and tree density of the area or community being mapped. As a result, without LiDAR (only available for the mainland portions of BISC) or stereo imagery, these parameters become very difficult to establish, thus further diminishing the potential of the computer-based classification to distinguish between communities whose spectral signature are similar or identical but differ only in canopy height (e.g. Red Mangrove Shrubland vs Red Mangrove Forest). As a result, for a project of this size, the logistics of using a computer-based classification was determined to be far less efficient and prone to a higher degree of error than the ocular identification and hand digitizing of vegetation communities.

#### *The Vegetation of Biscayne National Park:*

Ideally, mapping efforts should coincide with the acquisition of the imagery being used. In practice, however, this is rarely the case. Realistically because of delays associated in contracting and budgeting, and image acquisition, availability, and/or procurement, it is not uncommon, for most mapping efforts, to be initiated several months to a year or more after the imagery was initially flown. Moreover, the extent, scale, and detail required of a map along with the work force available can extend the temporal separation between the imagery and the final product by several more months or years (see Welch et al. 1999). Luckily, most vegetative communities tend to be relatively inert and resistant to structural and/or compositional change during the lifespan of most mapping projects. However, because of perturbations, both natural and anthropogenic, communities sometimes change rapidly creating conditions on the ground that are no longer coincident with the imagery. This has the unfortunate effect of creating a classification impasse for the cartographer who must decide between mapping what is currently there and known against what might have been prior to the perturbation event. This is also a major problem in the accuracy assessment phase of all mapping project.

For this project, the imagery dated to May 2005 (see methods section), but mapping efforts did not commence until nearly two years later, in March 2007. Consequently, the map (Plate 1) represents the vegetation communities of BISC between March 2007 and June 2008 when the first and last polygons were drawn, respectively, and when ground referencing and truthing efforts ceased, in February 2008. Arguably and reasonably, the point could also be made, based on the image acquisition date, that this map really represents the vegetation of BISC in

2005. However, since all mapping activities were conducted nearly three years after the imagery was taken, and the accuracy assessment won't be complete until nearly a year after map production, it is best to consider it a representation of the vegetation communities of BISC in 2007-08, or for brevity, 2008.

However, ongoing exotic removal and restoration efforts, particularly on Elliott Key and Boca Chita, created conditions on the ground that were no longer coincident with the imagery or definable within the classification system. As a result, on Elliott Key, for example, areas treated for *Thespesia populnea* (Seaside Mahoe) were systematically classified based on the 2005 spectral signature which suggested a monotypic community of this species. These areas could have been classified as Treated Seaside Mahoe. However, conditions on the ground, in 2008, were ambiguous and indicative of a transitional state between treated and non-treated Seaside Mahoe communities with some units completely devoid of vegetation and others showing signs of an emerging mixed mangrove Shrubland, dotted with emerging Seaside Mahoe seedling and saplings, in a succulent herbaceous matrix. Moreover, since the imagery did not allow for the delineation of these new transitional units we had no choice but to map them, by default, based on how they appeared on the imagery. Fortunately, this is an exception and not the norm within the map.

The map created for the project (Plate 1) encompasses a total area of 35.2 km<sup>2</sup> of which 31 km<sup>2</sup> are within the borders of BISC. Nearly 55% (17.0 km<sup>2</sup>) of these 31 km<sup>2</sup> correspond to the island habitat alone. The residual 4.2 km<sup>2</sup> (i.e. the difference between the total area mapped and the area mapped corresponding to BISC) consist of sections of the mainland that are adjacent to the park boundary (Figure 1). While not within the legislative jurisdiction of the National Park Service and BISC, these wetlands are an important component of the overall protection and ecological health of the communities that fall within the Park, including Biscayne Bay. As a result, some efforts were directed towards mapping these ecologically important lands. Unfortunately, only a small percentage of the total area in the BBCW were mapped.

Forest communities within BISC; i.e. Mangrove Forest & Hammock Forest, on both the mainland and islands, account for 48.4% of the total area mapped (Table 6). In contrast, the Shrubland and Scrub communities combine for almost 49% of all vegetation area (Table 6). The remaining area was distributed unevenly between Woodland (2%), Marsh (0.2%), Dune (0.03%), and Exotic (0.3%) communities (Table 6).

The total forested area within the Park was greater on the islands than on the mainland, 9.0 km<sup>2</sup> vs 5.7 km<sup>2</sup>, respectively (Table 6). However, mangrove forest communities were more abundant on the mainland than on the islands (Table 6). The Coastal Hardwood Hammock community with a total area of 7.023 km<sup>2</sup> (i.e. 22.7% of the total area mapped) was the overall dominant community type present within all of BISC (Table 6) but only occurred on the islands. Woodlands are a minor component of both island and mainland habitats. The distribution of Woodlands were slightly greater on the islands than in the mainland. Red and Mixed Mangrove Shrublands on the islands account for almost 3 times the total area of shrublands of any sort on the mainland (Table 6). In contrast, the scrub mangrove communities of the mainland accounted for almost 4 times the total area of scrub habitat of all

types on the islands (Table 6). Marshes, like Woodlands, were a minor component of both island and mainland habitats. However, Marshes were more common on the mainland than on the islands (Table 6). The Dune class was the only class found exclusively on the islands but it only accounted for 0.1% of the total area mapped within the island community. Exotics, at 0.27% of the total area mapped, were a minor component of both the mainland and island habitats.

Descriptively, the vegetation of BISC (Plate 1) consists of a complex mosaic of physiognomically distinct communities distributed along a hydrologic and salinity gradient that, along with nutrient availability and substrate type, ultimately determines plant community composition and productivity. The physiography of the landscape serves as a template controlling hydrology and salinity, and thereby regulating the distribution and structure of the communities. Soil type and depth are equally important, particularly on the mainland portions of BISC where the topographic gradient is subtle (Meeder et al. 2000) and soil depths can reach and exceed 1 m (Gaiser et al. 2006, Meeder et al. 2000). In contrast, soil depth on the upland portions of the islands rarely exceeds a few decimeters, but the topographic gradient between adjacent mangrove and hardwood hammock communities may easily exceed 1 meter. This juxtaposition in topography and substrate (both in type and depth) between the mainland and islands create unique environments that support unique communities within BISC (Table 6). This is particularly true for the Marsh class, where three of the four Level 3 Marsh types documented (e.g. Herbaceous Salt Marsh, Succulent Salt Marsh, and Graminoid Freshwater Prairie) are found exclusively in the island environment but not on the mainland (Table 6). There are other examples of this, most notably, the two Hammock Forest types that are exclusive to the islands (Table 6). Coincidentally, both of these hammock communities, though floristically and structurally similar, have distinctive soil characteristics, which sets them apart. Exclusive to the mainland, on the other hand, we find extensive areas of *Rhizophora mangle* dwarf trees (red mangrove scrub) in a matrix of either *Fimbristylis spadicosa* (Marsh Fimbry) or *Juncus roemerianus* (Black Rush), as well as isolated Salt Marsh communities dominated by Black Rush. It is worth noting that these Black Rush Salt Marshes are a relic community that is slowly being displaced by the encroachment of mangroves, particularly *R. mangle*.

Table 6: Summary statistics for all **Level – 3** communities, except Exotic (**Level – 1**) identified and mapped within BISC.

Class (L1)	Group (L3)	Area (km <sup>2</sup> )			Percent		
		Mainland	Islands	Total	Mainland	Islands	Total
Forest	Black Mangrove Forest	0.0571	0.399	0.457	0.4	2.4	1.47
	Buttonwood Forest	0.0000	0.081	0.081	0.0	0.5	0.26
	White Mangrove Forest	0.0000	0.000	0.000	0.0	0.0	0.00
	Red Mangrove Forest	0.4478	0.319	0.767	3.2	1.9	2.48
	Mixed Mangrove Forest	5.2444	1.110	6.354	37.3	6.6	20.53
	Coastal Hardwood Hammock	0.0000	7.023	7.023	0.0	41.6	22.69
	Coastal Dune Hammock	0.0000	0.034	0.034	0.0	0.2	0.11
Woodland	Black Mangrove Woodland	0.1343	0.185	0.319	1.0	1.1	1.03
	Buttonwood Woodland	0.0000	0.115	0.115	0.0	0.7	0.37
	White Mangrove Woodland	0.0000	0.042	0.042	0.0	0.3	0.14
	Mixed Mangrove Woodland	0.0229	0.097	0.120	0.2	0.6	0.39
	Upland Hardwood Woodland	0.0000	0.027	0.027	0.0	0.2	0.09
Shrubland	Black Mangrove Shrubland	0.0000	0.081	0.081	0.0	0.5	0.26
	Buttonwood Shrubland	0.0000	0.093	0.093	0.0	0.6	0.30
	White Mangrove Shrubland	0.0097	0.011	0.021	0.1	0.1	0.07
	Red Mangrove Shrubland	0.4085	2.525	2.934	2.9	14.9	9.48
	Mixed Mangrove Shrubland	1.6943	3.100	4.794	12.1	18.3	15.49
	Coastal Hardwood Shrubland	0.0000	0.063	0.063	0.0	0.4	0.20
Scrub	Black Mangrove Scrub	0.0000	0.101	0.101	0.0	0.6	0.33
	White Mangrove Scrub	0.0000	0.030	0.030	0.0	0.2	0.10
	Red Mangrove Scrub	4.9972	0.807	5.804	35.5	4.8	18.75
	Mixed Mangrove Scrub	0.5742	0.374	0.948	4.1	2.2	3.06
	Upland Scrub	0.0000	0.002	0.002	0.0	0.0	0.01
	Upland Hardwood Scrub	0.0000	0.015	0.015	0.0	0.1	0.05
Marsh	Graminoid Salt Marsh	0.0341	0.0005	0.035	0.2	0.0	0.11
	Herbaceous Salt Marsh	0.0000	0.001	0.001	0.0	0.0	0.00
	Succulent Salt Marsh	0.0000	0.004	0.004	0.0	0.0	0.01
	Graminoid Freshwater Prairie	0.0000	0.017	0.017	0.0	0.1	0.05
Dune	Mixed Herbaceous Dune	0.0000	0.009	0.009	0.0	0.1	0.03
Exotic	Exotic	0.0314	0.051	0.082	0.2	0.3	0.26
Other	Barren Microkarst	0.0000	0.020	0.020	0.0	0.1	0.06
	Barren Salt Flat	0.0000	0.004	0.004	0.0	0.0	0.01
	Beach	0.0000	0.022	0.022	0.0	0.1	0.07
	Lightning Gap	0.0013	0.0001	0.001	0.0	0.0	0.00
	Littoral Zone	0.0000	0.002	0.002	0.0	0.0	0.01
	Water	0.1738	0.040	0.214	1.2	0.2	0.69
	Anthropogenic	0.2248	0.097	0.322	1.6	0.6	1.04
<b>Total</b>		<b>14.1</b> <b>(45.4%)</b>	<b>16.9</b> <b>(54.6%)</b>	<b>31.0</b>	<b>100</b>		

## **Conclusion:**

By providing a spatial inventory of the plant communities within BISC, this map, directs managers to focus their attention on relic, rare, and fragmented communities (e.g. *J. roemerianus* salt marshes found on the mainland or mangrove shrublands in a matrix of *Spartina bakeri* found on to Elliott Key) that are generally more vulnerable to anthropogenic and natural perturbations. This is particularly true in the case of the species rich Coastal Hardwood Hammock community. This fragmented upland coastal community serves as refugia for many threatened and endangered species, both flora and fauna. However, since this community is only found on the eastern rim of Biscayne Bay, at a few meters above mean sea level, it is highly vulnerable to the effects of tropical storms and hurricanes and their accompanying storm surge, and, without question, sea-level rise. Moreover, in conjunction with existing data, managers can use this map to isolate communities that are likely to benefit the most from restoration efforts. Finally, this map, with its highly accurate and precise 1:300 scale shoreline, serves as a turn-of-the-21<sup>st</sup>-century baseline for the extent of mangroves within Biscayne National Park. As a result, it can be useful for monitoring the effects of sea-level rise on the wetlands and forested communities of Biscayne National Park.

## **Acknowledgements:**

This project was sponsored by the National Park Service South Florida / Caribbean Network Inventory and Monitoring Program under cooperative agreement H500 06 5040. Without question, this project could not have reach completion without the assistance and dedication of Kevin R. T. Whelan, Andrea Atkinson, Judd Patterson, Brian Witcher, & Matt Patterson from the sponsoring agency. Likewise, I would like to thank Steve L. Newman & Max Tritt from Biscayne National Park for their logistical support in providing much needed transportation and equipment essential to complete this project. Special thanks go out to the staff and pilots of HMC helicopters in particular Gary Freeman and Mark Palmieri for their hospitality and professionalism. I would also like to thank Mike Kline, Brooke Shamblin, and Pete Harlem along with Jonathan Moser, Jose Luciani, & Eric Sudalter, for assisting in the collection of ground-reference points. Pete Harlem was also instrumental in the unearthing of many historical documents as well as aerial photographs, which contributed to our understanding of the geology and ecology of Biscayne National Park. Lastly, we would like to thank Jay P. Sah, Keqi Zhang, Susana L. Stoffella, & Jack Meeder for their comments and support during this challenging but rewarding project.

## Literature Cited:

- Benz, U. C., P. Hoffmann, G. Willhauck, I. Lingenfelder, & M. Heynen. 2004. Multi-resolution, object-oriented fuzzy analysis of remote sensing data for GIS-ready information. *ISPRS Journal of Photogrammetry and Remote Sensing*. 58: 239-258.
- Bork, E. W., J. G. Su. 2007. Integrating LIDAR data and multispectral imagery for enhanced classification of rangeland vegetation: A meta analysis. *Remote Sensing of the Environment*. 111: 11-24.
- Davis, J. H., 1943, Vegetation map of Southern Florida 1:400,000. Bulletin 25: Tallahassee, Florida Geological Survey.
- Davis, J. H. 1967. General map of natural vegetation of Florida. 1:1,250,000. *Univ. of Florida*. Gainesville.
- Definiens Imaging. 2006. Definiens Professional User Guide 5. <http://www.definiens.com>.
- Gaiser, E. E., A. Zafiris, P. L. Ruiz, F. A. C. Tobias, & M. S. Ross. 2006. Tracking rates of ecotone migration due to salt-water encroachment using fossil mollusks in coastal South Florida. *In* J.C. Trexler, E. E. Gaiser, & D.L. Childers (eds), *Interaction of Hydrology and Nutrients in Controlling Ecosystem Function in Oligotrophic Coastal Environments of South Florida*. *Hydrobiologia*. 569: 237-257.
- Gaiser, E. E., A. Wachnicka, P. L. Ruiz, F. Tobias, & M. S. Ross. 2005. Diatom indicators of ecosystem change in subtropical coastal wetlands. *In* Bortone, S. A. (ed.) *Estuarine Indicators*. pp. 127-144. CRC Press, Boca Raton, FL, USA. 531 pp.
- Hill, R. A., & A. G. Thomson. 2005. Mapping woodland species composition and structure using airborne spectral and LiDAR data. *International Journal of Remote Sensing*. 26(17): 3763-3779.
- Hyypä, J., O. Kelle, M. Lehtikoinen, & M. Inkinen. 2001. A segmentation-based method to retrieve stem volume estimates from 3-D tree heights models produced by laser scanners. *IEEE Transaction on Geoscience and Remote Sensing*. 39(5): 969-975.
- IHRC. 2004. Windstorm simulation & modeling project: Airborne LiDAR data and digital elevations models in Miami-Dade, Florida. Final Report. Florida International University, International Hurricane Research Center.
- Lefsky, M. A., W. B. Cohen, & T. A. Spies. 2001. An evaluation of alternate remote sensing products for forest inventory, monitoring and mapping of Douglas fir forest in western Oregon. *Canadian Journal of Forestry Research*. 28: 1016-1031.

- Leynes, J B., and D. Cullison. 1998. Biscayne National Park historic resource study. National Park Service, Southeast Region. Atlanta, Georgia.
- Meeder, J. F., P. W. Harlem, M. S. Ross, E. E. Gaiser, & R. Jaffe. 2000. Southern Biscayne Bay Watershed Historic Creek Characterization: Tasks 1, 2, & 3. Report to the South Florida Water Management District.
- Ross, M. S., J. J. O'Brien, and L. L. Sternberg. 1994. Sea-level rise and the reduction in pine forests in the Florida Keys. *Ecological Applications*. 4(1): 144-156.
- Ross M. S. & P. L. Ruiz. 2003. Vegetation. *In*: E. E. Gaiser & M. S. Ross, Water Flow Through Coastal Wetlands. Annual Report to Everglades National Park. December 2003.
- Ruiz, P. L. and M. S. Ross. 2004. Hydrologic restoration of the Biscayne Bay Coastal Wetlands: mosquito and drainage ditch inventory and recommendations. Report to Biscayne National Park, Homestead, FL, USA.
- Rutchev, K., T. Schall, B. Doren, A. Atkinson, M. S. Ross, D. T. Jones, M. Madden, L. Vilchek, K. Bradley, J. Snyder, J. Burch, T. Pernas, B. Witcher, M. Pyne, R. White, T. J. Smith, J. Sadle, C. S. Smith, M. Patterson, and G. Gann. 2007. Vegetation Classification System for South Florida Natural Areas. Version 5.22.07.
- Stephenson, T. A. & A. Stephenson. 1950. Life between tide-marks in North America: The Florida Keys. *The Journal of Ecology*. 38(2): 354-402.
- Suárez, J. C., C. Ontiveros, S. Smith, and S. Snape. 2005. Use of airborne LiDAR and aerial photography in the estimation of individual tree heights in forestry. *Computers & Geoscience*. 31: 253-262.
- Welch, R., M. Madden, and R. F. Doren. 1999. Mapping the Everglades. *Photogrammetric Engineering & Remote Sensing*. 65(2): 163-170.
- Zhang, K., P.A. Houle, M.S. Ross, P.L. Ruiz, and M. Simard. 2006. Airborne laser mapping of mangroves on the Biscayne Bay coast, Miami, Florida. *Proceedings of 2006 IEEE International Geoscience and Remote Sensing Symposium & 27th Canadian Symposium on Remote Sensing*: 3750-3754, July 31-August 04, 2006, Denver, Colorado, USA.
- Zhang, K., M. Simard, M. S. Ross, V. H. Rivera-Monroy, A. Houle, P. L. Ruiz, R. R. Twilley, & K. R. T. Whelan. 2008.. Airborne laser scanning quantification of disturbances from hurricanes and lightning strikes to mangrove forest in Everglades National Park, USA. *Sensors*. 8: 2262-2292.

# **Plate 1**

**The 2008 Terrestrial Vegetation of Biscayne National Park**

# THE 2008 TERRESTRIAL VEGETATION OF BISCAYNE NATIONAL PARK



## Vegetation

### Forest

- Hammock Forest
- Coastal Hardwood Hammock
- Coastal Dune Hammock
- Black Mangrove Forest
- Buttonwood Forest
- White Mangrove Forest
- Red Mangrove Forest
- Mixed Mangrove Forest

### Woodland

- Upland Hardwood Woodland
- Black Mangrove Woodland
- Buttonwood Woodland
- White Mangrove Woodland
- Mixed Mangrove Woodland

### Shrubland

- Coastal Hardwood Shrubland
- Upland Shrubland
- Black Mangrove Shrubland
- Buttonwood Shrubland
- White Mangrove Shrubland
- Red Mangrove Shrubland
- Mixed Mangrove Shrubland

### Scrub

- Upland Scrub
- Upland Hardwood Scrub
- Black Mangrove Scrub
- White Mangrove Scrub
- Red Mangrove Scrub
- Mixed Mangrove Scrub

### Marsh

- Succulent Salt Marsh
- Herbaceous Salt Marsh
- Graminoid Salt Marsh
- Graminoid Freshwater Prairie
- Graminoid Freshwater Marsh

### Dune

- Mixed Herbaceous Dune


### Exotic

- Exotic
- Australian Pine
- Treated Australian Pine
- Seaside Mahoe

### Other

- Anthropogenic
- Barren Microkarst
- Barren Salt Flat
- Beach
- Lightning Gap
- Littoral Zone
- Inland Lakes or Pools
- Unclassified

Biscayne National Park Boundary



Southeast Environmental Research Center  
**SOUTH FLORIDA TERRESTRIAL ECOSYSTEMS LAB**

Florida International University, Miami, FL 33199

Authored by:  
 Funded by:  
 Date:  
 Base Map Source:  
 Datum:

Pablo L Ruiz  
 National Park Service South Florida / Caribbean Network  
 June 2008  
 2005 Multispectral (RGBNIRPan) Aerial Photos  
 NAD83 UTM Zone17N (units = meters)

# **Appendix 1**

## **Hierarchical Classification System with Community Descriptions**

## *L1 - Class*

<i>Class</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Forest</b> High-density stands of trees (> 50% tree canopy cover) with heights > 5 meters.	RutcheY et al.	100000	100000
<b>Woodland</b> Low-density stands of trees (10 - 60% tree canopy cover) with heights > 5 meters in a matrix of shrubs, graminoids, and/or herbaceous vegetation.	RutcheY et al.	200000	200000
<b>Shrubland</b> Stands of small trees and/or shrubs (canopy cover ≥ 50%) with heights < 5 meter tall.	RutcheY et al.	300000	300000
<b>Scrub</b> Communities of dwarf trees or shrubs typically in a matrix of graminoids, and/or herbaceous vegetation. Canopy cover 10% to 50% but can be as high as 100% for Mangrove and Cypress classes. Canopy < 5 meters tall with the exception being for Mangrove which is ≤ 2 meters.	RutcheY et al.	400000	400000
<b>Marsh</b> Graminoid and/or herbaceous emergent or floating vegetation in shallow water that stands at or above the ground surface for much of the year.	RutcheY et al.	500000	500000
<b>Dune</b> A ridge of wind blown or windstorm deposited sand or similar material directly inland and parallel to the shoreline that is commonly vegetated by graminoids and/or herbs and sometimes even shrubs.	RutcheY et al.	600000	600000
<b>Submerged Aquatic Vegetation</b> Vegetation that has evolved the ability to carry out its entire life cycle completely submerged in an aquatic environment.	RutcheY et al.	700000	700000
<b>Exotic</b> Non-native and invasive vegetation.	RutcheY et al.	800000	800000
<b>Other</b> Non-vegetative or anthropogenic cover.	Ruiz & Ross	900000	900000
<b>Unclassified</b> Unclassified vegetation or land cover.	Ruiz & Ross	999999	

## L2 - Type

<i>Type</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Mangrove Forest</b> Regularly flooded (tidal) forest found along coastal areas dominated by salt tolerant species.	RutcheY et al.	110000	110000
<b>Hammock Forest</b> Rarely inundated and well drained forests containing a mixture of tropical and temperate broad-leaved trees.	RutcheY et al.	130000	130000
<b>Mangrove Woodland</b> Regularly flooded (tidal) open canopy forest found along coastal areas dominated by salt tolerant species.	RutcheY et al.	210000	210000
<b>Upland Woodland</b> Rarely inundated and well drained open canopy forests containing a mixture of tropical and temperate broad-leaved trees.	RutcheY et al.	230000	230000
<b>Mangrove Shrubland</b> Regularly flooded (tidal) shrubland found along coastal areas dominated by salt tolerant species	RutcheY et al.	310000	310000
<b>Upland Shrubland</b> Rarely inundated and well drained shrublands containing a mixture of tropical and temperate broad-leaved trees.	RutcheY et al.	340000	340000
<b>Mangrove Scrub</b> Tidal and seasonally flooded dwarf (< 2 m height) mangrove trees found along coastal areas and in the transition zone between freshwater and marine dominated environments. Canopy densities are generally between 10% - 50% but can be as high as 100%.	RutcheY et al.	410000	410000
<b>Upland Scrub</b> Upland graminoid and/or herbaceous dominant communities in a matrix of dwarf (< 2 m height) trees and/or shrubs.	RutcheY et al.	430000	430000
<b>Salt Marsh</b> Marsh consisting of salt tolerant (halophilic) graminoid and/or herbaceous vegetation.	RutcheY et al.	510000	510000
<b>Freshwater Marsh</b> Marsh consisting of freshwater graminoid and/or herbaceous vegetation.	RutcheY et al.	520000	520000
<b>Herbaceous Dune</b> Herbaceous dominated dune.	RutcheY et al.	620000	620000

Appendix 1 - Hierarchical Classification System with Community Descriptions.

<i>L2 - Type</i>	<i>Author</i>	<i>Raster ID</i>	<i>Ruthey ID</i>
<b>Australian Pine</b> Macrophyte community consisting of River Sheoak ( <i>Casuarina cunninghamiana</i> ), Australian Pine ( <i>C. equisetifolia</i> ), and/or Suckering Australian Pine ( <i>C. glauca</i> ).	Ruthey et al.	803000	803000
<b>Treated Australian Pine</b> Macrophyte community treated for the presence of River Sheoak ( <i>Casuarina cunninghamiana</i> ), Australian Pine ( <i>C. equisetifolia</i> ), and/or Suckering Australian Pine ( <i>C. glauca</i> ).	Ruthey et al.	804000	804000
<b>Seaside Mahoe</b> Macrophyte community consisting of <i>Thespesia populnea</i> .	Ruthey et al.	833000	833000
<b>Non-Vegetative</b> Non-vegetative coverage	Ruiz & Ross	910000	
<b>Anthropogenic</b> Non-natural coverage associated with human infrastructure and/or activities.	Ruiz & Ross	920000	

## L3 - Group

<i>Group</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Black Mangrove Forest</b> Black Mangrove ( <i>Avicennia germinans</i> ) dominated forest found along the coast.	RutcheY et al.	111000	111000
<b>Buttonwood Forest</b> Buttonwood ( <i>Conocarpus erectus</i> ) dominated forest usually found on the landward edge of the coastal mangrove zone.	RutcheY et al.	112000	112000
<b>White Mangrove Forest</b> White Mangrove ( <i>Laguncularia racemosa</i> ) dominated forest found along the coast.	RutcheY et al.	113000	113000
<b>Red Mangrove Forest</b> Red Mangrove ( <i>Rhizophora mangle</i> ) dominated forest found along the coast.	RutcheY et al.	114000	114000
<b>Mixed Mangrove Forest</b> Mixed mangrove forest with no particular species of dominance found along the coast.	RutcheY et al.	115000	115000
<b>Coastal Hardwood Hammock</b> Forest containing a mixture of tropical and temperate broad-leaved trees found along coastal areas on rocky substrate overtoped by an organic layer.	RutcheY et al.	131000	131000
<b>Coastal Dune Hammock</b> Forest containing a mixture of tropical and temperate broad-leaved trees found along coastal areas on a dune (shell mounds excluded).	Ruiz & Ross	137000	
<b>Black Mangrove Woodland</b> Black Mangrove ( <i>Avicennia germinans</i> ) in a matrix composed of salt marsh graminoids, herbs, and/or succulents.	RutcheY et al.	211000	212000
<b>Buttonwood Woodland</b> Buttonwood ( <i>Conocarpus erectus</i> ) in a matrix composed of salt marsh graminoids, herbs, and/or succulents.	RutcheY et al.	212000	211000
<b>White Mangrove Woodland</b> White Mangrove ( <i>Laguncularia racemosa</i> ) in a matrix composed of salt marsh graminoids, herbs, and/or succulents.	Ruiz & Ross	213000	
<b>Mixed Mangrove Woodland</b> Mixed assemblage of mangrove tree species in a matrix composed of salt marsh graminoids, herbs, and/or succulents.	Ruiz & Ross	215000	

<i>L3 - Group</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Upland Hardwood Woodland</b> Woodland containing a mixture of tropical and temperate broad-leaved trees found along coastal areas on rocky substrate with little or no soil.	RutcheY et al.	233000	233000
<b>Black Mangrove Shrubland</b> Black Mangrove ( <i>Avicennia germinans</i> ) dominant shrubland predominately found in the upper part of the intertidal zone or on higher elevations.	RutcheY et al.	311000	311000
<b>Buttonwood Shrubland</b> Buttonwood ( <i>Conocarpus erectus</i> ) dominant shrubland usually found on the landward edge of the coastal mangrove zone or on the edge of hammocks.	RutcheY et al.	312000	313000
<b>White Mangrove Shrubland</b> White Mangrove ( <i>Laguncularia racemosa</i> ) dominant shrubland found throughout the intertidal zone.	RutcheY et al.	313000	314000
<b>Red Mangrove Shrubland</b> Red Mangrove ( <i>Rhizophora mangle</i> ) dominant shrubland found on the middle and lower portions of the intertidal and upper subtidal zone.	RutcheY et al.	314000	315000
<b>Mixed Mangrove Shrubland</b> Mixed of mangrove shrubland with no particular species of dominance found along the coast.	RutcheY et al.	315000	316000
<b>Coastal Hardwood Shrubland</b> Shrubland containing a mixture of tropical and temperate broad-leaved trees found along coastal areas on rocky substrate with little or no soil.	RutcheY et al.	342000	342000
<b>Black Mangrove Scrub</b> Black Mangrove ( <i>Avicennia germinans</i> ) dominant scrub predominately found in the upper part of the intertidal zone or on higher elevations.	RutcheY et al.	411000	411000
<b>White Mangrove Scrub</b> White Mangrove ( <i>Laguncularia racemosa</i> ) dominant scrub found throughout the intertidal zone.	RutcheY et al.	413000	413000
<b>Red Mangrove Scrub</b> Red Mangrove ( <i>Rhizophora mangle</i> ) dominant scrub found on the middle and lower portions of the intertidal and upper subtidal zone.	RutcheY et al.	414000	414000
<b>Mixed Mangrove Scrub</b> Mixed of mangrove scrub with no particular species of dominance found along the coast.	RutcheY et al.	415000	415000
<b>Upland Hardwood Scrub</b> Scrub containing a mixture of tropical and temperate broad-leaved trees found along coastal areas on rocky substrate with little or no soil.	RutcheY et al.	431000	431000

Appendix 1 - Hierarchical Classification System with Community Descriptions.

<i>L3 - Group</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Graminoid Salt Marsh</b> Graminoid dominated salt marsh	RutcheY et al.	511000	511000
<b>Herbaceous Salt Marsh</b> Herbaceous dominated salt marsh.	RutcheY et al.	512000	512000
<b>Succulent Salt Marsh</b> Succulent dominated salt marsh.	RutcheY et al.	514000	514000
<b>Graminoid Freshwater Marsh</b> Graminoid dominated freshwater marsh.	RutcheY et al.	522000	522000
<b>Graminoid Freshwater Prairie</b> Short hydroperiod freshwater marsh characterized by a mixture of low-stature grasses and sedges.	RutcheY et al.	523000	523000
<b>Mixed Herbaceous Dune</b> Mixed herbaceous dominated dune.	Ruiz & Ross	623000	
<b>Barren Salt Flat</b> Barren, generally hypersaline, flats exposed at low tide.	RutcheY et al.	910010	907000
<b>Beach</b> Sand and fine shell and coral fragments found along the shoreline.	RutcheY et al.	910020	901000
<b>Littoral Zone</b> Shoreline that is submerged at high tide and exposed at low tide and is usually devoid of upland vegetation.	Ruiz & Ross	910030	
<b>Lightning Gap</b> Canopy gaps created by cloud to ground lightning strikes.	Ruiz & Ross	910050	
<b>Mud</b> Moist or dry open ground.	RutcheY et al.	910070	903000
<b>Water</b> Open water areas such as ponds, lakes, rivers, bays, and estuaries.	RutcheY et al.	910120	904000
<b>Barren Microkarst</b> Karst topography devoid of vegetation usually found around the perimeter of Coastal Hardwood Hammocks.	Ruiz & Ross	910130	

Appendix 1 - Hierarchical Classification System with Community Descriptions.

<i><b>L3 - Group</b></i>	<i><b>Author</b></i>	<i><b>Raster ID</b></i>	<i><b>Rutchey ID</b></i>
<b>Campground</b> Area designated for camping	Ruiz & Ross	920010	
<b>Canal</b> Water bodies specifically designed to direct water from one location to another.	Rutchey et al.	920020	902020
<b>Dock</b> A place used as a landing place or moorage for boats.	Ruiz & Ross	920030	
<b>Lawns &amp; Landscaping</b> Ground that is covered with grass and/or trees and is maintained for its esthetics.	Ruiz & Ross	920040	
<b>Levee</b> Elevated berm, generally with an access road, utilized to contain a body of water such as a lake or marsh	Rutchey et al.	920050	902060
<b>Parking Lot</b> Area used for the parking of motor vehicles.	Ruiz & Ross	920060	
<b>Quarry</b> Area used for mining rocks, minerals, or other natural resources.	Rutchey et al.	920070	902080
<b>Road</b> Paved and unpaved roads other than levees.	Rutchey et al.	920080	902100
<b>Seawall</b> A wall or embankment to protect the shore from erosion or to act as a breakwater.	Ruiz & Ross	920090	
<b>Trail</b> A marked or established path or route designed to be followed.	Ruiz & Ross	920110	
<b>Spoil</b> Earth and rock excavated or dredged.	Rutchey et al.	920120	902110
<b>Walkway</b> A path, passage, etc. for pedestrians.	Ruiz & Ross	920120	
<b>Agriculture</b> Fields designated for the production of goods or food through the cultivation of plants.	Rutchey et al.	921000	902010

*Appendix 1 - Hierarchical Classification System with Community Descriptions.*

<i><b>L3 - Group</b></i>	<i><b>Author</b></i>	<i><b>Raster ID</b></i>	<i><b>Rutchey ID</b></i>
<b>Building</b> A roofed and walled structure built for permanent use.	Ruiz & Ross	922000	
<b>Unknown</b> Unknown land cover.	Rutchey et al.	929999	908000

## L4 - Formation

<i>Formation</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Black Mangrove-Buttonwood Forest</b> Black mangrove ( <i>Avicennia germinans</i> ) and Buttonwood ( <i>Conocarpus erectus</i> ) trees with the cover of either species ranging between 25-75%.	RutcheY et al.	115100	115100
<b>Black Mangrove-White Mangrove Forest</b> Black mangrove ( <i>Avicennia germinans</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) trees with the cover of either species ranging between 25-75%.	RutcheY et al.	115200	115200
<b>Black Mangrove-Red Mangrove Forest</b> Black mangrove ( <i>Avicennia germinans</i> ) and Red Mangrove ( <i>Rhizophora mangle</i> ) trees with the cover of either species ranging between 25-75%.	RutcheY et al.	115300	115300
<b>Buttonwood-White Mangrove Forest</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) trees with the cover of either species ranging between 25-75%.	RutcheY et al.	115400	115400
<b>Black Mangrove Woodland-Succulent</b> Black Mangrove ( <i>Avicennia germinans</i> ) trees in a matrix composed predominately of succulents.	RutcheY et al.	211030	211030
<b>Buttonwood Woodland-Succulent</b> Buttonwood ( <i>Conocarpus erectus</i> ) trees in a matrix composed predominately of succulents.	RutcheY et al.	212030	211030
<b>Buttonwood-White Mangrove Woodland</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) trees with the cover of either species ranging between 25-75%.	Ruiz & Ross	215400	
<b>Black Mangrove Shrubland-Succulent</b> Black Mangrove ( <i>Avicennia germinans</i> ) shrubs in a matrix composed predominately of succulents.	Ruiz & Ross	311040	
<b>Buttonwood Shrubland-Succulent</b> Buttonwood ( <i>Conocarpus erectus</i> ) shrubs in a matrix composed predominately of succulents.	Ruiz & Ross	312040	
<b>White Mangrove Shrubland-Succulent</b> White Mangrove ( <i>Laguncularia racemosa</i> ) shrubs in a matrix composed predominately of succulents.	Ruiz & Ross	313040	
<b>Mixed Mangrove Shrubland-Succulent</b> Mixed mangrove shrubs in a matrix composed predominately of succulents.	Ruiz & Ross	315040	

Appendix 1 - Hierarchical Classification System with Community Descriptions.

<i><b>LA -Formation</b></i>	<i><b>Author</b></i>	<i><b>Raster ID</b></i>	<i><b>Rutchey ID</b></i>
<b>Black Mangrove-White Mangrove Shrubland</b> Black mangrove ( <i>Avicennia germinans</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) shrubs with the cover of either species ranging between 25-75%.	Rutchey et al.	315200	316200
<b>Black Mangrove-Red Mangrove Shrubland</b> Black mangrove ( <i>Avicennia germinans</i> ) and Red Mangrove ( <i>Rhizophora mangle</i> ) shrubs with the cover of either species ranging between 25-75%.	Rutchey et al.	315300	316300
<b>Buttonwood-White Mangrove Shrubland</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) shrubs with the cover of either species ranging between 25-75%.	Rutchey et al.	315500	316500
<b>Buttonwood-Red Mangrove Shrubland</b> Buttonwood ( <i>Conocarpus erectus</i> ) and Red Mangrove ( <i>Rhizophora mangle</i> ) shrubs with the cover of either species ranging between 25-75%.	Rutchey et al.	315600	316600
<b>White Mangrove-Red Mangrove Shrubland</b> White Mangrove ( <i>Laguncularia racemosa</i> ) and Red Mangrove ( <i>Rhizophora mangle</i> ) shrubs with the cover of either species ranging between 25-75%.	Rutchey et al.	315700	316700
<b>Black Mangrove Scrub-Succulent</b> Black Mangrove ( <i>Avicennia germinans</i> ) scrub in a matrix composed predominately of succulents.	Rutchey et al.	411040	411040
<b>White Mangrove Scrub-Succulent</b> White Mangrove ( <i>Laguncularia racemosa</i> ) scrub in a matrix composed predominately of succulents.	Rutchey et al.	413040	413040
<b>Red Mangrove Scrub-Graminoid</b> Red Mangrove ( <i>Rhizophora mangle</i> ) scrub in a matrix composed predominately of graminoids.	Rutchey et al.	414010	414010
<b>Red Mangrove Scrub-Open Marsh</b> Red Mangrove ( <i>Rhizophora mangle</i> ) scrub in a matrix composed predominately of Open Marsh or Open Salt Marsh.	Rutchey et al.	414030	414030
<b>Red Mangrove Scrub-Dominant</b> Greater than 50% areal coverage of dwarf Red Mangrove ( <i>Rhizophora mangle</i> ) trees.	Rutchey et al.	414050	414050
<b>Mixed Mangrove Scrub-Graminoid</b> Mixed mangrove scrub in a matrix composed predominately of graminoids.	Rutchey et al.	415010	415010
<b>Mixed Mangrove Scrub-Succulent</b> Mixed mangrove scrub in a matrix composed predominately of succulents.	Rutchey et al.	415040	415040

Appendix 1 - Hierarchical Classification System with Community Descriptions.

<i><b>LA -Formation</b></i>	<i><b>Author</b></i>	<i><b>Raster ID</b></i>	<i><b>RutcheY ID</b></i>
<b>Black Mangrove-White Mangrove Scrub</b> Black mangrove ( <i>Avicennia germinans</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees with the cover of either species ranging between 25-75%.	RutcheY et al.	415200	415200
<b>Black Mangrove-Red Mangrove Scrub</b> Black mangrove ( <i>Avicennia germinans</i> ) and Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees with the cover of either species ranging between 25-75%.	RutcheY et al.	415300	415300
<b>Buttonwood-White Mangrove Scrub</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees with the cover of either species ranging between 25-75%.	RutcheY et al.	415400	415400
<b>Buttonwood-Red Mangrove Scrub</b> Buttonwood ( <i>Conocarpus erectus</i> ) and Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees with the cover of either species ranging between 25-75%.	RutcheY et al.	415500	415500
<b>White Mangrove-Red Mangrove Scrub</b> White Mangrove ( <i>Laguncularia racemosa</i> ) and Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees with the cover of either species ranging between 25-75%.	RutcheY et al.	415700	415600
<b>Black Rush</b> Black Rush ( <i>Juncus roemerianus</i> ) dominated salt marsh.	RutcheY et al.	511200	511200
<b>Cordgrass</b> Sand Cordgrass ( <i>Spartina bakeri</i> ) and/or Gulf Cordgrass ( <i>S. spartinae</i> ) dominated salt marsh.	RutcheY et al.	511400	511400
<b>Glasswort</b> Glasswort ( <i>Salicornia spp.</i> ) dominated salt marsh.	RutcheY et al.	514200	514200
<b>Sawgrass</b> Sawgrass ( <i>Cladium jamaicense</i> ) dominated marsh.	RutcheY et al.	522100	522100
<b>Commercial</b> A building or complex housing retail business.	Ruiz & Ross	922010	
<b>Government</b> A building or complex housing government offices.	Ruiz & Ross	922020	
<b>Historical</b> A building or complex with historical significances	Ruiz & Ross	922030	

*Appendix 1 - Hierarchical Classification System with Community Descriptions.*

<b><i>L4 -Formation</i></b>	<b><i>Author</i></b>	<b><i>Raster ID</i></b>	<b><i>Rutchey ID</i></b>
<b>Industrial</b> A building or complex housing factories.	Ruiz & Ross	922040	
<b>Lighthouse</b> A tower or structure designed to emit light and aid in navigation.	Ruiz & Ross	922050	
<b>Pump Station</b> A Structure used to move water through canals.	Rutchey et al.	922060	902040
<b>Residential</b> A building or complex used as a permanent dwelling.	Ruiz & Ross	922070	

## L5 - Alliance

<i>Alliance</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Black Mangrove Woodland-Saltwort</b> Black Mangrove ( <i>Avicennia germinans</i> ) trees in a matrix composed predominately of Saltwort ( <i>Batis maritima</i> ).	Ruiz & Ross	211031	
<b>Buttonwood-White Mangrove Woodland-Herbaceous</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) trees in a matrix composed predominately of herbaceous vegetation.	Ruiz & Ross	215420	
<b>Black Mangrove Shrubland-Saltwort</b> Black Mangrove ( <i>Avicennia germinans</i> ) shrubs in a matrix composed predominately of Saltwort ( <i>Batis maritima</i> ).	Ruiz & Ross	311041	
<b>Buttonwood Shrubland-Saltwort</b> Buttonwood ( <i>Conocarpus erectus</i> ) shrubs in a matrix composed predominately of Saltwort ( <i>Batis maritima</i> ).	Ruiz & Ross	312041	
<b>Black Mangrove-White Mangrove Shrubland-Succulent</b> Black Mangrove ( <i>Avicennia germinans</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) shrubs in a matrix composed predominately of succulent vegetation.	Ruiz & Ross	315210	
<b>Buttonwood-White Mangrove Shrubland-Succulent</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) shrubs in a matrix composed predominately of succulent vegetation.	Ruiz & Ross	315510	
<b>Buttonwood-White Mangrove Shrubland-Graminoid</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) shrubs in a matrix composed predominately of graminoids.	Ruiz & Ross	315520	
<b>Black Mangrove Scrub-Saltwort</b> Black Mangrove ( <i>Avicennia germinans</i> ) dwarf trees in a matrix composed predominately of Saltwort ( <i>Batis maritima</i> ).	RutcheY et al.	411041	411041
<b>White Mangrove Scrub-Saltwort</b> White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees in a matrix composed predominately of Saltwort ( <i>Batis maritima</i> ).	RutcheY et al.	413041	413041
<b>White Mangrove Scrub-Glasswort</b> White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees in a matrix composed predominately of Glasswort ( <i>Salicornia bigelovii</i> ).	RutcheY et al.	413042	413042
<b>Red Mangrove Scrub-Sawgrass</b> Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees in a matrix composed predominately of Sawgrass ( <i>Cladium jamaicense</i> ).	RutcheY et al.	414011	414011

<i>L5 -Alliance</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Red Mangrove Scrub-Frimbry</b> Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees in a matrix composed predominately of Marsh Frimbry ( <i>Fimbristylis spadicea</i> ).	RutcheY et al.	414014	414014
<b>Red Mangrove Scrub-Black Rush</b> Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees in a matrix composed predominately of Black Rush ( <i>Juncus roemerianus</i> ).	RutcheY et al.	414015	414015
<b>Red Mangrove Scrub-Subtidal</b> Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees in a subtidal environment.	Ruiz & Ross	414051	
<b>Mixed Mangrove Scrub-Black Rush</b> Mixed mangrove dwarf trees in a matrix composed predominately of Black Rush ( <i>Juncus roemerianus</i> ).	RutcheY et al.	415015	
<b>Black Mangrove-White Mangrove Scrub-Succulent</b> Black Mangrove ( <i>Avicennia germinans</i> ) & White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees in a matrix composed predominately of succulents.	RutcheY et al.	415240	415240
<b>Buttonwood-White Mangrove Scrub-Graminoid</b> Buttonwood ( <i>Conocarpus erectus</i> ) & White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees in a matrix composed predominately of graminoids.	RutcheY et al.	415410	415410
<b>Buttonwood-White Mangrove Scrub-Succulent</b> Buttonwood ( <i>Conocarpus erectus</i> ) & White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees in a matrix composed predominately of succulents.	RutcheY et al.	415440	415440
<b>Buttonwood-Red Mangrove Scrub-Graminoid</b> Buttonwood ( <i>Conocarpus erectus</i> ) & Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees in a matrix composed predominately of graminoids.	RutcheY et al.	415510	415510

## L6 - Association

<i>Association</i>	<i>Author</i>	<i>Raster ID</i>	<i>RutcheY ID</i>
<b>Buttonwood-White Mangrove Shrubland-Cordgrass</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) shrubs in a matrix composed predominately of Sand Cordgrass ( <i>Spartina bakeri</i> ).	Ruiz & Ross	315521	
<b>Black Mangrove-White Mangrove Scrub-Saltwort</b> Black Mangrove ( <i>Avicennia germinans</i> ) -White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees in a matrix composed predominately of Saltwort ( <i>Batis maritima</i> ).	RutcheY et al.	415241	415241
<b>Buttonwood-White Mangrove Scrub-Cordgrass</b> Buttonwood ( <i>Conocarpus erectus</i> ) and White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees in a matrix composed predominately of Sand Cordgrass ( <i>Spartina bakeri</i> ).	RutcheY et al.	415417	415417
<b>Buttonwood-White Mangrove Scrub-Glasswort</b> Buttonwood ( <i>Conocarpus erectus</i> ) & White Mangrove ( <i>Laguncularia racemosa</i> ) dwarf trees in a matrix composed predominately of Glasswort ( <i>Salicornia bigelovii</i> ).	RutcheY et al.	415442	415442
<b>Buttonwood-Red Mangrove Scrub-Sawgrass</b> Buttonwood ( <i>Conocarpus erectus</i> ) & Red Mangrove ( <i>Rhizophora mangle</i> ) dwarf trees in a matrix composed predominately of Sawgrass ( <i>Cladium jamaicense</i> ).	RutcheY et al.	415511	415511

## **Appendix 2**

**Definiens Professional® v5 Segmentation Algorithm**

In *Definiens Professional®* v5, two process trees were created and saved – one for coastal shorelines and one for island shorelines. They are named Land\_Water\_Boundary\_Coast and Land\_Water\_Boundary\_Keys. Both processes contain the same steps, but parameters were altered within the listed functions. Within the Process Tree, processes are grouped into the general areas of Segmentation, Classification, Merge objects and Export. The Process Tree for the Keys is shown in Fig. 1.



Fig. 1: Process Tree for shoreline creation using *Definiens Professional®* V5. This tree is used for the Keys. The Process Tree used for coastal areas is similar.

To create a project in *Definiens Professional®* v5, we loaded the image of interest, then defined the No Data areas. The near infrared band rarely reads zero in the scene; therefore we use band 4 = 0 as the definition of the No Data area. Next, image objects were created using multiresolution segmentation. In multiresolution segmentation, there are several parameters that can be evaluated. For both the keys and the coastal area, the best segmentation results were seen using only the NIR band 4 of the image and adjusting the Composition of Homogeneity Criteria to 0.6 for color and 0.4 for shape. This ratio refers to the relative importance of spectral input and shape for the image objects. The shape criteria was placed as high as 0.4 to give image objects that were more compact and therefore easier to edit later in ArcGIS as a shapefile.

Fig. 2a & 2b show the segmentation parameters for the coastal area and the keys. Because the coastal area is simpler in vegetation patterns, the scale parameter is 150 rather than 75 in the keys. The scale parameter is used as a threshold for heterogeneity calculations to determine if two objects are similar enough to be merged or kept separate. The remaining parameters are the same in the two multiresolution segmentation processes.




Fig. 2a: Multiresolution segmentation parameters for coastal area.




Fig. 2b: Multiresolution segmentation parameters for keys.

Classification of the image objects is defined through the Class Hierarchy. Fig. 3 shows the Class Hierarchy. There are two levels in the hierarchy. In the first level, objects are classified as Water or Other (not water). The Water class has an explicit definition; whereas the Other class is defined as those objects not classed as Water. In the second level, the Other objects are further classified into Wetter Area and Land (not wetter area). Again, here the Wetter Area class has an explicit definition, and the Land Class are those objects not classed as Wetter Area.




Fig. 3: Class Hierarchy for classification of objects as Land or Water.

To help reduce the influence of the spectral variability of the images in the classification, a ratio, NDVI, was defined as a custom feature using the brightness values:

$$NDVI = (NIR - Red)/(NIR + Red) = (Band 4 - Band 1)/(Band 4 + Band 1)$$

Classes were then classified as followed (Fig 4.)


Coastal Area

The 'Class Description' dialog for 'Water' in the 'Coastal Area' shows the name 'Water' with a blue color swatch. The parent class is 'Water'. Under the 'All' tab, the 'Contained' section shows a logical expression: 'and (min) Mean Layer 1 <= 90 NDVI < 0.1'.



Keys

The 'Class Description' dialog for 'Water' in the 'Keys' area shows the name 'Water' with a blue color swatch. The parent class is 'Water'. Under the 'All' tab, the 'Contained' section shows a logical expression: 'and (min) NDVI <= -0.1'.

### Coastal Area



### Keys



The Wetter Area class includes areas having shallow water and NDVI just above the cutoff for deeper water. For the coastal area the Wetter Area class is defined as a threshold using NIR (Band 4  $\leq$  50).




Fig 4: class descriptions and functions used for classification.

In contrast the definition of Wetter Area of the keys is defined by a fuzzy less than function with a mean value of 95.

In general the differences between the keys and the coastal areas are:

Water is defined by a lower threshold for the keys ( $\leq -0.1$  vs.  $< 0.1$ ).

Band 1 (Red) is also included in the coastal area.

Wetter Area is defined in the NIR as a threshold  $\leq 50$  for the coastal area and fuzzy less than 90-100 for the keys.

These class descriptions, which are included with the process tree and rule set, are loaded for each analysis. However, since each scene is different in its spectral properties, it was necessary to alter the rules to get adequate delineation of the shoreline.

It was also necessary to manually change object boundaries and classifications due to the four factors described there. There are a couple of ways to manually assign classes to objects. To get all of the features needed, click on the Manual Editing Toolbar Button (Fig. 5) rather than using the Process Tree. The lower limit for resolving mangrove tree islands was a diameter of about 10 meters, depending on the quality of the photograph.




Fig. 5.: Manual Editing Toolbar.

When the objects and classes closely follow the shorelines in the image, then objects can be merged and exported. In V4 (Ecognition), classification based image object fusion could be done in a couple of steps. In V5 the following steps must be done to merge adjoining objects having the same class:

1. Copy all the objects and their classification to a 'higher level'. The image object fusion will be done at this new level while preserving the original segmentation and classification.
2. Use the Merge Region function for each class individually. This is real important, otherwise the entire image will end up as a single large object. The Merge Classes steps in the Process Tree are shown in Fig. 6, and the individual steps are shown in Fig. 7.



Fig. 6.: Merging objects portion of Process Tree.

Note: The processes nested under each category are read from bottom to top.





Fig. 7.: Parameters for copying level and merging objects by class.

At this point, the image has been segmented and the objects classified as Land, Wetter Areas (shallow water) and Water (Deep water). For the purposes of creating shorelines, the project is simplified by changing the class of Wetter Areas to Water then repeating the object fusion. Assigning objects to a particular class is done through the Process Tree (Fig. 8).




Fig. 8.: Assign class to an object. Here objects classed as Wetter Area are changed to the Water class.

The result of the classification in Fig 9. The green area are objects classified as Land, and the blue areas are classified as water.




Fig. 9: Results of classification based on algorithm developed.

The last step in the process is to export the objects as a shapefile. The shapefile can be assigned a projection and edited in ArcGIS.