

**Landscape Pattern–Marl Prairie/Slough Gradient; patterns and trends in Shark Slough
marshes and associated marl prairies**

(Cooperative Agreement #: W912HZ-14-2-0023)

Annual Report: Year 2 (2015/2016)



Submitted to

Ms. Sherry Whitaker

U.S. Army Engineer Research and Development Center (U.S. Army - ERDC)

3909 Halls Ferry Road, Vicksburg, MS 39081-6199

Email: Sherry.L.Whitaker@usace.army.mil

Jay P. Sah, Michael S. Ross, Susana Stoffella,

Rosario Vidales, Alexander Martinez-Held

Southeast Environmental Research Center,

Florida International University,

Miami, FL 33186

2017

Authors' Affiliation

Jay P. Sah, Ph.D. – *Research Associate Professor*

Florida International University
Southeast Environmental Research Center
11200 SW 8th Street, Miami, FL 33199
Tel. (305) 348-1658
sahj@fiu.edu

Michael S. Ross, Ph.D. – *Professor*

Florida International University
Southeast Environmental Research Center/Department of Earth & Environment
11200 SW 8th Street, Miami, FL 33199
Tel. (305) 348-1420
rossm@fiu.edu

Susana Stoffella, M.S. – *Research Analyst*

Florida International University
Southeast Environmental Research Center
11200 SW 8th Street, Miami, FL 33199
Tel. (305) 348-0493
stoffell@fiu.edu

Rosario Vidales – *Research Technician*

Florida International University
Southeast Environmental Research Center
11200 SW 8th Street, Miami, FL 33199
Tel. (305) 348-6066
rvida018@fiu.edu

Alexander Martinez-Held – *Research Technician*

Florida International University
Southeast Environmental Research Center
11200 SW 8th Street, Miami, FL 33199
Tel. (305) 348-6066
aheld@fiu.edu

Table of Contents

Authors' Affiliation	ii
Table of Contents	iii
General Background	1
1. Introduction	2
2. Methods	3
2.1 Study Sites	3
2.2 Vegetation sampling	5
2.3 Water depth measurements	5
2.5 Data Analysis	6
2.4.1 Hydroperiod and annual mean water depth	6
2.4.2 Fire frequency and Time since last fire	6
2.4.3 Vegetation classification and ordination	7
2.4.4 Biomass estimation	7
2.4.5 Vegetation response to hydrology – Trajectory analysis	7
2.4.6 Weighted averaging and Vegetation-inferred hydroperiod	8
2.4.7 Soil-hydrology relationship	8
3. Results	8
Soil characteristics	13
4. Discussion & Conclusion	26
5. References	29

General Background

The Monitoring and Assessment Plan (MAP), the primary tool of the REstoration COordination and VERification (RECOVER) program to assess the performance of the Comprehensive Everglades Restoration Plan (CERP) provides the data and analytical support necessary to implement adaptive management. In the Everglades, marsh vegetation in both marl prairie and ridge and slough landscapes is sensitive to large-scale restoration activities associated with the CERP authorized by the Water Resources Development Act (WRDA) of 2000. More specifically, changes in hydrologic regimes at both local and landscape scales are likely to affect vegetation composition in the transition zone between these two landscapes, resulting in a shift in boundary between plant communities. In order to track these dynamics, Florida International University (Dr Michael Ross-PI and Dr. Jay Sah-Co-PI) has undertaken a study of vegetation structure and composition in relation to physical and biological processes along the marl prairie-slough gradient since 2005. Since the Fall of 2014, the study has been led by Dr. Jay Sah, while Dr. Michael Ross is also actively involved as the Co-PI in the study.

Vegetation monitoring transects in the Shark Slough basin, funded by US Army Corps of Engineers (USACOE) under RECOVER-MAP, capture the full range of marl prairie and slough plant communities, and address Performance Measure (PM):GE-15(Landscape Pattern–Marl Prairie/Slough gradient),by “...detecting spatio-temporal change in vegetation structure and composition in response to natural and restoration-induced hydrologic changes...”. Monitoring of vegetation along the marl prairie/slough gradients addresses a working hypothesis that ‘Spatial patterning and topographic relief of ridges and sloughs are directly related to the volume, timing and distribution of sheet flow and related water depth patterns’, identified in the hypothesis cluster “Landscape Patterns of Ridge and Slough Peatlands and Adjacent Marl Prairies in Relation to Sheet Flow, Water Depth Patterns and Eutrophication” (RECOVER 2009). The study also addresses the hypothesis that, with the implementation of CERP, resumption of historical flow and related patterns of hydroperiod, water depth, and fire will cause a noticeable change in plant community composition and structure along the gradient and in the transition zone between marl prairie and peat-dominated ridge and sloughs.

Initiated in 2005 as an expansion on Shark Slough study transects that had been established and sampled in 1998-2000 with funding from DOI’s Critical Ecosystems Study Initiative (CESI), the ongoing study is in the US Army Corps of Engineers (USACE)-funded fourth sampling cycle beginning from Fall 2014. This document summarizes results for Year-2 (2015/2016) of the current cycle (2014-2018), and it updates the previous year (Year-1 of the current cycle) results. It now includes a summary of vegetation dynamics along Transect M3, which was initially sampled in 2006/2007, and then sampled every three years (2009/2010, 2012/2013, and 2015/2016).

1. Introduction

In the Everglades, plant communities arranged along environmental gradients are expressions of ecosystem functional processes associated with underlying physico-chemical drivers that vary in space and time. Hence, determining the responses to spatio-temporal changes in key environmental drivers of plant assemblages along gradients, and the boundaries between them, is important for conservation and ecosystem restoration. The landscapes in both Shark River and Taylor Slough basins of the Everglades include long hydroperiod sloughs, flanked by short hydroperiod marl prairies. Particularly in the Shark River Slough (SRS) basin, vegetation structure and composition change gradually along an elevation and water depth gradient, from short-hydroperiod marl prairies to the ridge and slough landforms that epitomize central SRS (Olmsted and Loope 1984; Olmsted and Armentano 1997; Ross et al. 2003). Hydrology is one of the major drivers of species differences between marl prairie and ridge-and-slough landscapes. Hence, alterations in hydrologic conditions usually cause a shift in vegetation structure and composition within each landscape; extreme changes can lead to even dominance of hydric vegetation in marl prairie or various levels of degradation of landforms in the ridge and slough (R&S) landscape. In the past century, changes in the amount and flow patterns of water, resulting from the construction and operation of a series of canals, levees and water structures (Light and Dineen 1994, McVoy et al. 2011), have altered the proportions of prairie and slough vegetation in the region. Furthermore, changes in water management associated with the ongoing Comprehensive Everglades Restoration Plan (CERP 2000) are likely to affect vegetation composition in the transition zone between these ecosystems, resulting in a shift in the boundary between marl prairie and slough communities. It is therefore important to understand how restoration impacts the dynamics of prairie and slough landscapes and the boundaries between the two.

Along the gradient from marl prairies to ridge-and-slough marshes, vegetation in the prairie is likely to respond more rapidly to hydrologic changes than vegetation in the marshes. Armentano et al. (2006) argued that the transition from one vegetation type to another (e.g., prairie to marsh) in response to hydrology may take place in as little as 3 to 4 years. However, the transition from marsh to prairie appears to take longer. In the southern Everglades, recent water management efforts have been directed at ameliorating the adverse effects of previous water management activities. In this respect, a series of water detention ponds have been brought into operation along the eastern boundary of the Park to mitigate the wet-season water reversals that were prevalent in this region due to loss of water from the rocky glades to the canal (Van Lent et al. 1999). In contrast, strategic regulation of water deliveries through the S12 structures along US 41 has been in place since 2002, with the intent of reversing damage that was caused by the extended wet conditions that resulted from both high water deliveries and rains in the mid-to-late 1990s. These modifications in water management activities, along with those planned under the Central Everglades Planning Project (CEPP), including the construction and operation of two Tamiami Trail bridges, have affected and will continue to affect water conditions within the Park, resulting in changes in vegetation communities and ecological processes.

In 2005, we initiated a long-term study of vegetation dynamics in relation to changes in underlying environmental drivers, especially hydrology, along the MP-S gradient. The broader goal of the study is to assess the impact of Everglades restoration activities on plant communities along the

gradient, and to detect any shifts in position and attributes of boundaries between those communities. The study is conducted on five transects that extend across SRS into adjacent marl prairies. Shark Slough portions of transects overlap with transects that were established and sampled under different sponsorship in 1998-2000, providing the prospect of assessing long-term temporal change in vegetation in those areas.

In this study, our specific objectives were, i) to characterize recent vegetation composition along the marl prairie-slough gradient, and ii) to assess changes in vegetation in both the Shark Slough and marl prairie portions of the transects over a seventeen-year period (1999-2016). We hypothesized that variation in vegetation composition along the MP-S gradient is driven primarily by hydrology, i.e., the duration and depth of flooding. We also hypothesized that Shark River Slough vegetation follows the temporal trend in hydrologic regimes, causing species composition to move toward assemblages more indicative of relatively dry conditions during the last fifteen years. In addition, in compliance with differential water management goals on the east and west sides of SRS, we hypothesized that marl prairie vegetation would follow the spatially differentiated temporal trend in hydrologic regimes. We expected vegetation in the eastern portion of the marl prairie to change toward a wetter character during 2006-2016, and vegetation in the western prairies to shift toward a drier type.

2. Methods

2.1 Study Sites

Sites sampled in 2015-2016 were part of the ongoing long-term vegetation monitoring along the marl prairie-slough gradient in the southern Everglades. The study design includes five transects (MAP Transect M1 to M5), varying in length from 9.0 km to 35.8 km. These transects were established in 2005, when the systematic sampling began along the gradient. Three transects, M1, M3 and M4 extend across the Shark Slough to adjacent short-hydroperiod marl prairie habitat (**Figure 1**). Transect M1, located in Northeastern Shark Slough (NESRS), extends only to the marl prairie on the east of the slough. Transect M2 originally covered an area restricted to Shark River Slough (SRS), extending on both sides of L-67S canal. But in 2015, this transect was extended eastward another 5 km, thereby covering prairie vegetation along the eastern boundary of the ENP and the transitional zone between the marl prairie in NESRS and ridge & slough landscape in SRS. Transect M5 covers an area between fresh to brackish water ecosystems in the southeastern corner of SRS, extending to the east into fresh water marl prairies located on both sides of the main Park road.

In Transect M3, the longest transect measuring 35.8 km, vegetation sampling plots were established at 200-500 m intervals. In the marl prairie section of the transect, the plots were established at 300 m intervals, and in the SRS portion of the transects, the plot density varied between 2 to 4 plots per km (200-500 meter intervals). Vegetation sampling on Transect M3 began in 2006, and the transect has been sampled every three years thereafter. **Table 1** summarizes the years and numbers of sites sampled on Transects M1-M3. The slough portion of these transects was sampled in the wet season (July to November), accessing the sites by airboat or helicopter. Marl prairie portions of the transect were sampled in the dry season (Jan to May) and were accessed by helicopter for drop off and pickup, and on foot for sampling. In 2016, however, dry season

sampling continued through June 13th, primarily due to unusual high water level in the early dry season, and the resulting difficulty in obtaining helicopter transport due to the high demand later in the spring by various research groups.




Figure 1: Location map of Marl prairie-Slough Gradient Study plots on Transects M1-M5.

Table 1: Sites sampled on MAP transects M3 between 2006 and 2016.

Transect	Sampling Event	Sites Sampled			
		Prairie sites		Slough sites	
		Year	No. of Sites	Year	Number of Sites
M1	E1	2006	11	2005	20
	E2	2009	11	2008	20
	E3	2012	11	2011	20
	E4	2015	11	2014	20
M2	E1			2005	25
	E2			2008	26
	E3			2011	25
	E4	2015	15	2014	25
M3	E1	2007	72	2006	37
	E2	2010	72	2009	37
	E3	2013	72	2012	37
	E4	2016	71	2015	37

2.2 Vegetation sampling

Vegetation was sampled in a nested-plot design that allowed for efficient sampling of the range of plant growth forms (herbs, shrubs and trees) present along the transects (Ross et al. 2005; Sah et al. 2013). Plots were sampled at 200-500m intervals. Higher intensity sampling occurred in areas accessible by airboat, based on the contention that increased sampling intensity would enable us to make a more meaningful comparison of current vegetation with that present on the same transects in 1999 (Ross et al. 2001; Ross et al. 2003).

At each sampling site, a PVC tube marked the SE corner of a 10x10m tree plot. Nested within each tree plot, a 5x5m herb/shrub plot was established, leaving a 1-m buffer strip along the southern and eastern border of the tree plot. In the 10x10m tree plots, we measured the DBH and crown length and width of any woody individual ≥ 5 cm DBH, and then calculated canopy cover assuming horizontally-flattened elliptical crown form. Within each 5x5m herb/shrub plot, we estimated the cover class of each species of shrub (woody stems >1 m height and <5 cm DBH) and woody vines, using the following categories: $<1\%$, 1-4%, 4-16%, 16-33%, 33-66%, and $>66\%$. We estimated the cover % of herb layer species (all herbs, and woody plants <1 m height) in five 1-m² subplots located at the four corners (NE, NW, SE and SW) and the center (CN) of the 5x5m plot. Species present in the 5x5m plot but not found in any of the 1m² subplots was assigned a mean cover of 0.01%. In addition, a suite of structural parameters was recorded in a 0.25m² quadrat in the SW corner of each of the five subplots. Structural measurements included the following attributes: 1) The height and species of the tallest plant in the plot; 2) Canopy height, i.e., the tallest vegetation present within a cylinder of ~ 5 cm width, measured at 4 points in each 0.25m² quadrat; 3) Total vegetative cover, in %, and 4) Live vegetation percent cover, expressed as a % of total cover.

2.3 Water depth measurements

Field water depths in combination with EDEN (Everglades Depth Estimation Network, <http://sofia.usgs.gov/eden>) water surface elevation data serve as the basis for calculation of ground elevation and estimation of hydrologic conditions at each site. Water depth was measured at each site along the transect, whether marl prairie or slough. We measured water depths at the PVC, the marker of the plot, and in the center of five vegetation sub-plots in a 5x5m plot. At the marl prairie sites of M1 and M3, water depths measured on a single day in the Fall of 2008, when there was standing water, were used to estimate hydrologic conditions.

2.4 Soil and Plant analysis

In 2015-2016, soil and plant samples were collected from 15 sites on Transect 3. Five sites were in the eastern portion of the marl prairie, five in the transition zone, and another five sites were within the ridge & slough portion of the transect. At each site, three of five 1x1m² sub-plots used for vegetation sampling were randomly selected for soil sampling. In each of these subplots, one soil core of 10 cm length was collected using 5.7 cm diameter core tube. At one site, M3-11400, soil samples were collected only in two sub-plots, as the other plots were very rocky and lacked enough soil to justify collection. Soil samples were placed in Ziploc bags, labelled, and brought to the lab at FIU, where they were refrigerated until further analysis. Each core's compaction and length were measured in the field and recorded.

The net weight of wet soil samples was obtained. Samples were oven-dried at 80°C for 48 hours or until a constant weight was achieved. Dry weight and volume were obtained, and the sample bulk density was calculated. We removed extraneous macro-materials, including roots and rocks. Samples were then ground to pass a 2-mm sieve. Later, 2-3 mg dry samples were obtained and enclosed in tin (Sn) capsules, and then delivered to the SERC Nutrient Analysis Lab at FIU for Total Nitrogen (TN) and Total Carbon (TC) Analysis. Another 2-3 g of dry sample were obtained and transferred to small screw cap glass vials (7 mL), and then delivered to FIU Freshwater Biochemistry Lab for Total Phosphorus (TP) and other analysis. Subsamples were also analyzed for Phosphatase and Glucosidase enzyme activity.

Plant samples were collected from a 0.25x0.25 m² quadrat within the same 1x1m² sub-plots in which soil samples were collected. Plant samples were then separated by species, oven dried at 70° C for 72 hrs, and weighed to calculate the above ground plant biomass. Plant samples for four major species, sawgrass (*Cladium jamaicense*), muhly grass (*Muhlenbergia capillaris* ssp. *filipes*), spikerush (*Eleocharis cellulosa*) and beak rush (*Rhynchospora tracyii*), were sub-sampled. Sawgrass was present in most of plots along the gradient, whereas muhly grass was restricted to the marl prairie portion of the gradient. Beakrush was common in marl prairie and the transition zone, while spikerush was present in the transition zone and ridge & slough portion of the gradient. The subsamples of these species were ground in a coffee-grinder. After each use, the coffee grinder was washed and dried to avoid contamination. The powdered samples were then sieved with a mesh (# 10 sieve), weighed, and placed into 20 mL scintillation glass vials for further analysis. Plant samples were stored for determination of total carbon (TC), total nitrogen (TN), $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$.

2.5 Data Analysis

2.4.1 Hydroperiod and annual mean water depth

We used field water depth-derived elevation and EDEN water surface elevation data to estimate the hydrologic conditions at each sampling site. We calculated the ground elevation of each plot using mean water depth for the plot and EDEN estimates of water surface elevation at that point (center of the plot) for the same sampling date. Daily water levels for each plot were estimated based on ground elevation and the time series data of water surface elevation extracted from the EDEN database. The hydroperiod, i.e., the number of days per year when a location had water depth >0cm, and mean annual water depth were calculated for each plot. We then averaged hydroperiod and mean annual water depth for the four water years (May^{1st}–April 30th) prior to each sampling event to examine vegetation response to hydrologic changes.

2.4.2 Fire frequency and Time since last fire

The fire geodatabase, in which the records of fire events are catalogued from 1948 to 2012, was obtained from Everglades National Park (ENP). The shape files for 2013-2016 fires were also obtained from the Park, and were later added to the geodatabase. The database contains shape files of fires with other attributes such as type of fire (Natural, RX, incendiary, etc.), date of incidence, etc. The data were used to calculate fire frequency and time since last fire (TSLF) for vegetation monitoring sites along Transect 3 using ArcGIS 10.3.

2.4.3 Vegetation classification and ordination

We summarized species data by calculating the importance value (IV) of each species present in herb and shrub layers in each plot. We calculated species' importance value as: $IV = (\text{relative cover} + \text{relative frequency})/2$. For calculating IV of the species that did not occur in any of five subplots but occurred in 5 x 5 m² plot, a frequency of 4% was assigned, on the assumption that such a species would have occurred in at least one subplot, had all 25 1 x 1 m² subplots within a plot been sampled. Vegetation types at all sites that were surveyed along the five transects between 2005 and 2008 had already been defined using a hierarchical agglomerative cluster analysis (Sah et al. 2013). In the analysis, Bray-Curtis dissimilarity was used as the distance measure, and the flexible beta method to calculate relatedness among groups and/or individual sites (McCune and Grace 2002). In the current year analysis, non-metric multidimensional scaling ordination (NMDS) was done to analyze the shift in species composition using trajectory analysis (see below sub-section 2.4.5)

2.4.4 Biomass estimation

For the sites in the marl prairie portion of the gradient, vegetation structural measurements were summarized for each plot, and mean canopy height and total vegetative cover were used to estimate above ground plant biomass, using the allometric equation developed by Sah et al. (2007) for marl prairie vegetation within CSSS habitat. The equation for calculating biomass was as follows:

$$\sqrt{\text{Biomass}} = 6.708 + 15.607 * \arcsin \sqrt{\text{Cover\%} / 100} + 0.095 * \text{Ht}$$

where Biomass = Total plant biomass (g/m²), Cover = Crown cover (%), and Ht = Mean crown height (cm).

2.4.5 Vegetation response to hydrology – Trajectory analysis

At the slough sites on Transects M1-M3, changes in vegetation composition since 1999/2000 were analyzed using trajectory analysis (Minchin et al. 2005; Sah et al. 2014), an ordination-based technique designed to test hypotheses about rates and directions of community change. In the NMDS ordination performed for trajectory analysis, we included vegetation data for prairie sites collected on all transects beginning with the first sampling cycle (2005-2008), and for the ridge and slough sites, we included all data collected between 1999 and 2016. Prairies sites were included to cover the full range of hydrologic conditions on the transects. In the NMDS ordination, the hydrology vector represented by mean annual water depth was defined through a vector fitting technique in DECODA (Kantvilas and Minchin 1989; Minchin 1998; Sah et al. 2014).

To quantify the degree and rate of change in vegetation composition along the reference vector, two statistics, delta (Δ) and slope were calculated (Minchin et al. 2005). Delta measures the total amount of change in the target direction. It was calculated as the difference between the projected score at the final time step and the initial time. Slope measures the mean rate of change in community composition along the target vector. The statistical significance of both delta (Δ) and slope was tested using Monte Carlo simulations with 10,000 permutations of the cover scores of

species among sampling times within each trajectory, with the NMDS ordination and calculation of trajectory statistics repeated on each permuted data matrix.

2.4.6 Weighted averaging and Vegetation-inferred hydroperiod

Vegetation change analysis in the marl prairie portion of the gradient included calculation of vegetation-inferred hydroperiod, i.e., the hydroperiod for a site indicated from its vegetation composition, using a weighted averaging partial least-square (WA-PLS) regression model. The training-data set with which we developed the WA-PLS regression model was the species cover data plus hydroperiod estimates from 291 plots on six topographically-surveyed transects within the Cape Sable seaside sparrow habitat (Ross et al. 2006). In developing the WA-PLS models, species cover were fourth square root transformed, which down-weights the influence of very dominant species. Mean hydroperiod was calculated over five years prior to vegetation sampling. The performance of the models was judged by the improvement in R^2 value and RMSEP (root mean square error of prediction). RMSEP was estimated by a leave-one-out (jackknife) cross-validation procedure, in which a vegetation-hydroperiod model is developed from all samples except one, and consequently applied to predict the hydroperiod of the left-out point on the basis of its vegetation. We used the C2 program of Juggins (2003) to develop the WA-PLS model.

Finally, the best WA-PLS model was applied to the calibration data set, here the MP-S gradient data that included vegetation data at 107 sites on Transect M3. The predicted hydroperiods for those sites were termed ‘vegetation-inferred hydroperiod’. A change in vegetation-inferred hydroperiod between successive sampling episodes reflects the amount and direction of change in vegetation, expressed in units of days (0-365) along a gradient in hydroperiod.

2.4.7 Soil-hydrology relationship

Soil nutrient data were summarized by sites and a person correlation was use to examine the relationships between hydrologic variables (hydroperiod and mean annual water depth) and soil characteristics.

3. Results

Hydrologic pattern (1999-2016)

In the slough portion of transects M1, M2 and M3, the hydroperiod and mean annual water depth averaged over four years prior to vegetation sampling varied over the study period. In the late 1990s, i.e. four years before the 1999 (E0) vegetation sampling, mean hydroperiod on all three transects were >360 days, and mean (\pm SD) annual water depths were 32.0 ± 10.7 , 44.2 ± 8.8 and 42.8 ± 10.3 cm on Transects M1, M2 and M3, respectively (**Figure 2**). At the slough sites on those transects, mean hydroperiod and annual water depth were lower during three subsequent sampling events (2005/2006 (E1), 2008/2009 (E2) and 2011/2012 (E3)) than during the initial; period. However, during 2014/2015 (E4) sampling, sites in the slough were wetter than the sampling done three years ago (**Figure 2**). In 2014/2015, four-year average hydroperiod at the slough sites on M1, M2 and M3 were 315, 337 and 338 days, and annual mean water depths were

25, 35 and 31 cm, respectively. Nonetheless, both hydroperiod and mean annual water depth were lower than late 1990s. The hydroperiod was 22-42 days shorter and mean water depth 9-12 cm less than the years before 1999 sampling.




Figure 2: Box Plots showing (A) hydroperiod and (B) mean annual water depth averaged over four years prior to vegetation sampling in the Shark Slough portions of MAP transects M1, M2 & M3. Different letters represent significant (pairwise t-test; $p < 0.05$) difference in (A) hydroperiod, and (B) mean annual water depth among years on individual transects.

Hydrologic conditions in the marl prairie portion of the transect M1 varied among sampling events. Mean hydroperiod, averaged over four years before E2 census was 60 days shorter than in the years before E1 (**Figure 3**). The hydrologic conditions in subsequent years, i.e. after E2 became wetter. The mean 4-year average hydroperiod before E3 was 18 days longer than the hydroperiod before E2, and the trend continued during the sampling event E4. In contrast to the hydroperiod, 4-year average annual mean water level was lowest during E3 (**Figure 4**). However, both the hydroperiod and mean annual water level before 2015 spring sampling (E4) was higher than E2 and E3, but was still shorter or less than the years before E1. The differences in hydrologic conditions between sampling events were mostly due to extreme events. The prolonged dry period between 2006 and 2008, i.e. the period before the 2nd census (E2) saw water levels dip far below the ground level (**Figure 5**). The water level in the spring of 2011, i.e. just before the 3rd census (E3) was also much lower than ground level. In the southern portion of NESRS, where Transect M2 was extended in 2015, four-year average hydroperiod was 292 ± 55 days (**Figure 3**), the value ranged from 161 days at the easternmost site to 346 days at the west end of the extended transect (M2E). Likewise, the four-year average annual mean water depth was 21.6 ± 17.8 cm, and it ranged between -15.5 cm to 45 cm.




Figure 3: Mean ($\pm 95\%$ CI) hydroperiod averaged over four years prior to vegetation sampling in the marl prairie portions of MAP transects M1, M2E, and M3. Different letters represent significant (pairwise t-test; $p < 0.05$) difference in hydroperiod between sampling period on individual transects or a section of the transect.




Figure 4: Mean ($\pm 95\%$ CI) annual water depth averaged over four years prior to vegetation sampling in the marl prairie portions of MAP transects M1, M2E and M3. Transect M3 has marl prairies both sides (East & West) of Shark Slough. Different letters represent significant (pairwise t-test; $p < 0.05$) difference in hydroperiod between sampling period on individual transects or a section of the transect

Transect M3 is unique, as the hydrologic conditions in the marl prairie portion of the transect differ between eastern and western marl prairies. Across all the sampling events, water condition was wetter in the eastern prairies than the western prairies. However, in the prairies on both sides of the slough, it was much drier during E2 than E1. In contrast, an increasing trend in both four-year average hydroperiod and mean annual water depth was observed during the next two sampling events, E3 and E4 (**Figure 3, 4**). On the both sides of the slough, the four-year average hydroperiod

showed its highest value during the last sampling year 2016 (**Figure 3**), but mean annual water depth was highest in 2016 only in eastern sites (**Figure 4**). It was important to note that, at the western prairie sites, despite unusually high water conditions in spring of 2016, the four-year mean annual water depth associated with the 2016 sampling was still significantly lower than during 2007 (E1) sampling. In 2006, the four-year average hydroperiod in eastern and western prairies were 260 ± 41 days and 224 ± 59 days, respectively. Likewise, the mean annual water depths were 9.3 ± 13.3 cm and 1.9 ± 10.4 cm, respectively.




Figure 5: 30-day average mean annual water depth on the marl prairie portion of the transect M1.

Fire frequency and time since last fire

Several sites on the MP-S gradient transects have burned frequently in the past. Between 1990 and 2005, the period that included vegetation sampling (1999/2000) at the slough sites, there were not many fires within the area. Nevertheless, since 2005, when vegetation monitoring began at regular interval on the transects, both prairie and slough sites on Transects M1, M2 & M3 burned in prescribed burns (Rx), human-caused fire or wildfires (**Table 2**). Time elapsed between the burned-year and sampling events, defined as time since last fire (TSLF), have affected vegetation composition observed at these sites.

Table 2: Vegetation sampling sites burned over the sampling period (2005-2015). The fire attributes were obtained from the Fire database of Everglades National Park.

Fire Name	Year	M1	M2	M3
L67 Rx	2005	0	1	0
Airboat	2006	18	4	7
U Road Rx	2007	0	10	0
Coptic	2007	1	0	0
West L67 WFU	2007	0	1	0
Mustang Corner	2008	11	1	44
Shark Valley Tram Rx	2009	0	0	1
ROG NE Rx	2012	0	12	31
EE 1 Rx	2012	18	13	0

Soil characteristics

Surface soil characteristics vary along MP-S gradient represented by the Transect M3. In general, soil total carbon (TC), organic carbon (OC) and total nitrogen (TN) showed an increasing trend from marl prairie to slough portions of the transect (**Figure 6a, b**). The mean (\pm) TC content ranged from $138.1 \pm 15.4 \text{ mg g}^{-1}$ at a prairie site near the ENP boundary to $444.1 \pm 15.1 \text{ mg g}^{-1}$ at a site (M3-18300) in the slough portion of the transect (**Table 3**). The mean OC ranged from $22.7 \pm 5.58 \text{ mg g}^{-1}$ to $404.0 \pm 13.2 \text{ mg g}^{-1}$, and mean TN varied between $2.79 \pm 0.99 \text{ mg g}^{-1}$ and $32.54 \pm 0.43 \text{ mg g}^{-1}$.

Table 3: Mean (± 1 SD) soil properties at the selected sites along MP-S gradient on the Transect M3.

Site ID	TN (mg g-1)	TC (mg g-1)	OC (mg g-1)	IC (mg g-1)	TP ($\mu\text{g g-1}$)
M3-00300	6.7 ± 0.66	138.1 ± 15.4	44.9 ± 2.8	93.2 ± 14.3	208.9 ± 34
M3-01500	6.1 ± 2.51	147.7 ± 13.9	43.5 ± 20.1	104.1 ± 7	163.4 ± 54.8
M3-02700	2.8 ± 0.99	138.8 ± 5.3	22.7 ± 5.6	116.2 ± 2.5	100.5 ± 20.9
M3-04800	7.3 ± 1.9	155.2 ± 4.4	51.9 ± 8.9	103.3 ± 5.7	197.6 ± 73
M3-06000	9.2 ± 2.74	141.9 ± 9.5	72.4 ± 23.8	69.5 ± 30.5	197.8 ± 35.8
M3-07200	13.4 ± 2.14	150.8 ± 14.9	138.8 ± 20.6	12 ± 16.4	337.8 ± 104.3
M3-08400	14.1 ± 5.47	170.3 ± 61.9	164.9 ± 60.8	5.4 ± 8.2	310 ± 111.2
M3-09900	7.6 ± 4.73	157.2 ± 19.8	81.5 ± 19.2	75.8 ± 0.5	261.9 ± 191.8
M3-11400	18.9 ± 10.08	270.8 ± 59.9	209.7 ± 114.9	61.1 ± 55	357.7 ± 164.4
M3-14000	10.6 ± 2.3	226.9 ± 28.6	119.5 ± 32.9	107.4 ± 4.3	177.3 ± 24.3
M3-16000	21.8 ± 12.84	321.4 ± 173.1	292.5 ± 176.9	28.9 ± 5.8	347.1 ± 25.5
M3-17000	28.1 ± 2.11	408.5 ± 17.8	381 ± 9.1	27.4 ± 9	269.9 ± 16.9
M3-18300	32.5 ± 0.43	444.1 ± 9.1	392.2 ± 19.2	51.8 ± 24.4	262.1 ± 25.1
M3-20800	28.5 ± 2.05	441.3 ± 10.1	399 ± 23.8	42.2 ± 16.3	367.4 ± 56
M3-22500	31.9 ± 1.31	408.5 ± 15.1	404 ± 13	4.5 ± 2.7	339 ± 29.2




Figure 6: Soil (a) total carbon (TC), organic carbon (OC, and (b) Total nitrogen (TN) content at sites along marl prairie-slough gradient on Transect M3

Unlike TC and OC, the mean IC showed a decreasing trend from prairie to slough. Soil IC ranged from the maximum of $116.2 \pm 2.5 \text{ mg g}^{-1}$ at a prairie site to $4.5 \pm 2.7 \text{ mg g}^{-1}$ at a slough site (**Table 3**). Soil total phosphorous (TP) did not show strong trend. However, in general, the prairie sites had lower soil phosphorous than the ridge and slough sites (**Figure 7**), but the variability is very high in the transition zone (**Table 3**). Mean (\pm SD) TP ranged between $100.5 \pm 20.9 \text{ } \mu\text{g g}^{-1}$ and $367.4 \pm 25.1 \text{ } \mu\text{g g}^{-1}$.




Figure 7: Soil total phosphorus (TP) content at sites along marl prairie-slough gradient on Transect M3.

Along the MP-S gradient, TC, OC and TN, were strongly and positively correlated ($r > 0.7$; $p = 0.001$) with hydroperiod and mean annual water depth, both averaged over 16 years (**Figure 8a-f**). Soil TP also increased with the two hydrologic variables. However, the correlation between soil TP and both variables were not significant in this study (**Figure 8g, h**). Unlike the TC, OC, TN and TP, soil IC decreased with the wetness of the sites, though the relationship was weak.




Figure 8: Relationships between hydrologic variables and soil TC, OC, TN and TP at the sites along marl prairie-slough gradient on the transect M3.

Shark River Slough vegetation change (1999-2015)

Between 1999 and 2015, marsh vegetation on all three transects (M1-M3), showed a shift in relative abundance of species indicative of sensitivity to the hydrologic changes. However, the direction and rate of vegetation change was not uniform throughout the sampling period. Trajectory analysis revealed that in the slough portion of M1 and M2 sampled at 3-6 year intervals between 1999 and 2014, species composition continued to shift towards drier vegetation types until 2011 (**Figures 9, 10**), but between 2011 and 2014, species composition shifted in the opposite direction, i.e. towards wetter vegetation types. In the slough portion of M3, the direction of vegetation shift was similar to that observed on M1 and M2 through 2009 (**Figure 11**). However,

during the next two sampling periods, its vegetation trajectory was opposite in direction to those observed on M1 and M2; between 2009 and 2012, the vegetation on M3 slough sites shifted towards a wetter type, whereas between 2012 and 2015 the sites showed a drying trend.




Figure 9: NMDS ordination bi-plots of site scores, the environmental vectors fitted in the ordination space, and the trajectory of centroid. The ordination is based on species abundance data collected five times between 1999 and 2014 in the Shark Slough portion of the Transect M1. Only the sites that showed significant ($p \leq 0.1$) rate of change in species composition along the hydrology gradient are shown. Initial point and the end of the trajectory represent the 1999 and 2014 sampling event, respectively

The percentage of sites that showed a drying or wetting trend in vegetation varied among transects, and in different portions of Transect 2. When summarized over the sampling period (1999-2015), most of sites during the E4 (2004/2015) sampling still showed drier vegetation types when compared to the conditions in 1999. Trajectory analysis revealed that the percent of sites with a significant shift towards dry vegetation was higher on M1 (50.0%) and M3 (53.6%) than M2 (33.3%), and east of the L67 levee than west of the levee on M2. On all three transects, the shift towards drier vegetation was greatest between the first two sampling events, E0 (1999) and E1 (2005). However, during subsequent periods, the vegetation change pattern differed among transects. Between E1 and E2 samplings, the shift towards dry vegetation continued on M1 and M3. In contrast, sites on M2 did not show a significant shift along the hydrologic gradient during that period (**Figure 10**).




Figure 10: NMDS ordination bi-plots of site scores, the environmental vectors fitted in the ordination space, and the trajectory of centroid. The ordination is based on species abundance data collected in the Shark Slough portion of the Transect M2. Only the sites that showed significant ($p \leq 0.1$) rate of change in species composition along the hydrology gradient are shown. Initial point and the end of the trajectory represent the 1999 and 2014 sampling event, respectively.




Figure 11: NMDS ordination bi-plots of site scores, the environmental vectors fitted in the ordination space, and the trajectory of centroid. The ordination is based on species abundance data collected in the Shark Slough portion of the Transect M3. Only the sites that showed significant ($p \leq 0.1$) rate of change in species composition along the hydrology gradient are shown. Initial point and the end of the trajectory represent the 1999 and 2015 sampling event, respectively.

Between E3 (2011/2012) and E4 (2014/2015), significant changes in species richness and relative abundance of some major species accompanied the trajectories described above. Mean species richness in the slough portion of M1 in 2014 was significantly higher than in 2011, but was similar to that in other sampling years (**Figure 12**). In the slough portion of M2, mean species richness in 2014 was significantly higher than any previous sampling years, except in 2005. During the E4 sampling, mean species richness at the M3 sites was also higher than other sampling years, but the difference was significant only between E0 and E4.




Figure 12: Plant species richness (species/plot) at the sites in slough portion of the transects M1, M2 and M3.

When averaged over all three transects, the total plant cover at the slough sites did not change much. However, relative abundance (Importance Value) of some of most abundant species (Mean Importance Value >2.0) changed significantly. In the previous sampling years, the drying trend was supplemented by an increase in relative abundance of sawgrass (*Cladium mariscus* ssp. *jamaicense*) and spikerush (*Eleocharis cellulosa*) and a decrease in abundance of bladderworts (*Utricularia* spp.) (**Figure 13**). However, during the latest sampling period (2011-2015), opposite trend was observed. Relative abundance of sawgrass significantly (Paired t-test; $p < 0.001$) decreased from 48.5% in 2011/2012 to 41.4% in 2014/2015 (**Figure 13a**). The mean abundance of spikerush did not differ between E3 and E4 samplings (**Figure 13b**). In contrast, within the

same period the relative abundance of eastern purple bladderwort (*Utricularia purpurea*) doubled, from 9.5% to 18.3% (**Figure 13c**). Interestingly, mean relative abundance of the other species of bladderwort (*Utricularia foliosa*) increased during the same period. As in previous sampling years, changes in the relative abundance of another two major species, lemon bacopa (*Bacopa caroliniana*) and maidencane (*Panicum hemitomom*), were not significant (**Figure 13e, f**).




Figure 13: Box-plots of major species' importance value (IV) in the slough portion of transects, averaged across all three M1, M2 & M3 transects for each sampling period.

Vegetation change in Marl Prairie (2005-2016)

In contrast to the vegetation change pattern observed in slough portion of the transect M1, trajectory analysis revealed that between 2012 and 2015 sampling events, species composition on prairie portion of this transect continued to shift towards a drier type (**Figure 14**). Likewise, more than 50% of sites showed that vegetation-inferred hydroperiod decreased. However, the magnitude of such changes in vegetation-inferred hydroperiod was much less (<30 days) than what was observed between 2006 and 2012 (>30 days) (**Figure 15**). Instead, during the latest sampling event there were more sites showing an increase in inferred hydroperiod than the sites before 2012. When averaged over all the sites, the mean reduction in inferred hydroperiod between 2006 and 2012 was 20 days, whereas between 2012 and 2015, it was only one day. Continued drying trend of some sites on M1 observed in 2014, resulted in a significant (t-test, $p < 0.05$) increase in relative abundance (IV) of Muhly grass (*Muhlenbergia capillaris* ssp. *filipes*), *Centella asiatica*, but a decrease in relative abundance of sawgrass and spikerush (Appendix 1). In contrast, relative abundance of some hydric species, such as Beakrush, (*Rhynchospora tracyi*) and Red Bacopa (*Bacopa caroliniana*) increased during the same period.




Figure 14: NMDS ordination bi-plots of the trajectory of centroid and the environmental vectors fitted in the ordination space. The ordination is based on species abundance data collected four times between 2006 and 2015 in the prairie portion of the Transect M1. Initial point and the end of the trajectory represent the 2006 and 2015 sampling event, respectively




Figure 15: Change in vegetation-inferred hydroperiod between different samplings at the vegetation monitoring plots on the marl prairie portion of the Transect M1.

On M3, which has marl prairie sites located both sides of SRS, vegetation change pattern differed between eastern and western prairies. While species composition in western prairies shifted towards a drier type, as evidenced in an increase in the abundance of *Schizachyrium rhizomatum*, the direction of change in vegetation composition in eastern prairie sites showed a mixed pattern (**Appendix 1**). Several sites at the distal portions of the transect, especially close to the eastern Park boundary, exhibited an increase in inferred-hydroperiod, suggesting that species composition at these sites shifted toward a wetter type between 2007 and 2016 (**Figure 16**). However, between the E3 (2013) and E4 (2016) samplings, the magnitude of change toward

wetter vegetation type along the eastern boundary of ENP was less than the shift over 10 year sampling period, between E1 (2007) and E4 (2016). Surprisingly, in recent years, the abundance of representatives of both prairie and hydric species were observed to increase on this section of the transect. Mean relative abundance of both sawgrass (*Cladium mariscus* ssp. *jamaicense*) and muhly grass (*Muhlenbergia capillaris* ssp. *filipes*) doubled in three years, from 2013 and 2016. The relative abundance of these species in 2016 was similar to the values in 2007.




Figure 16: Change in vegetation-inferred hydroperiod between different samplings at the vegetation monitoring plots on the marl prairie portion of the Transect M1.

The shift in vegetation composition observed over 10 years at the marl prairie portion M1 and M3 transects resulted also in changes in species richness and plant biomass. Mean species richness increased significantly (Pairwise t-test) on M1 and the eastern portion of M3W, where a drying trend was observed over 13 years. In contrast, on the western portion of M3E, species richness was significantly lower during the E4 than during any previous sampling years (**Figure 17**).




Figure 17: Plant species richness (species/plot) at the sites in marl prairie portion of the transects M1, M2E and M3. The Transect M3 extends within the marl prairies both sides (M3E and M3W) of Shark River Slough.

Mean plant biomass did not change on M1 (**Figure 18**), but did change on the eastern portion of M3 (M3E), where all but three plots burned in Mustang Fire in 2008. In the eastern marl prairie, biomass during the 2nd sampling (E2), two years after the fire, was only half of what it was during E1. Mean (\pm SD) aboveground biomass during E1 and E2 was $768 \pm 332 \text{ g m}^{-2}$ and $403 \pm 197 \text{ g m}^{-2}$, respectively. In this portion of M3, biomass recovered in three years, but at the time of E3, it was still only two-thirds of the initial biomass. In next three years, biomass increased slightly, but the mean biomass during E4 sampling ($541 \pm 273 \text{ g m}^{-2}$) was not significantly different (pairwise t-test; $df = 38$, $p = 0.428$) from the biomass during E3 ($514 \pm 267 \text{ g m}^{-2}$).




Figure 18: Above ground plant biomass (g m^{-2}) at the sites in marl prairie portion of the transects M1, M2E and M3. The Transect M3 extends within the marl prairies both sides (M3E and M3W) of Shark River Slough.

4. Discussion & Conclusion

In the Greater Everglades, the relationship between hydrologic regime and vegetation distribution is dynamic. Along the marl prairie-slough gradient, vegetation shift on individual transects was also influenced by year-to-year variation in water conditions, possibly caused by both rainfall and water management activities. For instance, while water level was above the thirty-year average during the mid-to-late 1990s, and continuously for three years prior to the 2005 sampling in northern SRS, water level was at or below the 30-year benchmark during the next four years. Moreover, in subsequent years, including the 2011 drought, mean annual water level varied greatly. However, the four-year average water level before 2014 sampling was higher than previous two (2008 & 2011) samplings (**Figure 2**). In concurrence with the hydrologic shift during the last sampling, vegetation composition in slough portions of M1 and M2 shifted toward a more hydric type (**Figures 8, 9**). In contrast, vegetation change on M3 was towards a drier type. Since not all transects were sampled in the same year, the annual variation in water conditions might have also affected the magnitude and direction of vegetation change on these transects. The sensitivity of vegetation change patterns to short-term variation in hydrologic conditions observed in this study supports earlier findings that in Everglades prairies and marshes, discernible change in species composition can occur in periods as short as 3-4 years (Armentano et al. 2006; Zweig and Kitchens 2008; Sah et al. 2014).

In the Everglades, the relative abundance of sawgrass and other hydric species such as spikerush, bladderwort and water lilies are considered as an indicator of water conditions in ridge and slough landscape (Ross et al. 2003; Zweig and Kitchens 2008; Ross et al. 2016). In this study, mean sawgrass cover showed an increasing trend until the 2012 sampling, while its cover decreased in next three years (**Figure 13**). In contrast, mean cover of bladderworts showed an opposite trend. The increase in sawgrass cover during the last 16 years in SRS supports the longer-term dynamics, described for the post-drainage era in the Everglades by Bernhardt and Willard (2009). Other researchers also have reported expansion of sawgrass and other emergent species, such as spikerush, in the R&S landscape, primarily due to decrease in water levels (Busch et al., 1998; Zweig and Kitchens, 2008, 2009; Nungesser 2011) and flow velocities (Larsen et al. 2011). Such expansion may occur within 3-4 years, especially when a minimum water level is maintained beneath the peat surface of the sloughs for three consecutive dry seasons (Zweig and Kitchens 2009). While the extensive expansion of sawgrass could be a step towards succession toward woody vegetation, especially when it occurs on elevated ground that experiences prolonged dry conditions, the extended wet seasons that occur intermittently in some years would reverse the process. In the slough portion of the transects M1, M2 and M3, both four-year average hydroperiod and mean annual water depth before E4 sampling was higher than before E3 (**Figure 2**). Results suggest that an increase of even 5-10 cm mean annual water depth in SRS can rapidly shift the vegetation towards one more characteristic of slough, as was evidenced in overall decrease in sawgrass and increase in bladderworts over three years, between E3 and E4 sampling events.

The deviation in trajectories of vegetation shift observed in the slough portion of transects is also affected by fire. Several sites on Transects M1-M3 burned between 2006 and 2012. The Mustang Corner fire that occurred in May 2008, following almost two years of drought and at a time when water level was 65 cm below the surface (Ruiz et al. 2013), may have consumed significant amounts of peat on the SRS portion of Transect M1. The vegetation at five burned sites on M1,

where the mean cover was 32% in 1999, was very sparse (cover 11.2%) and comprised mostly of hydric species during 2011 sampling. However, over the next three years, vegetation cover recovered, and mean species richness increased. Fire-induced elevation loss may also have contributed to compositional shift toward wetter vegetation at several locations on this Transect. In comparison to dry season fire, wet season fire seems to have less impact on the vegetation cover. All 18 slough sites on M1, and 25 sites on M2 were burned in two different prescribed fires in 2012 (Table 2), but the mean total cover in 2014, two and half years after fire, was already 65% of pre-burn cover.

Short-hydroperiod marl prairies in the Everglades are flooded annually for varying periods, while remaining dry for extended portions of the year. Generally, in seasonally-flooded ecosystems similar to the Everglades marl prairies, differences in optimum flooding tolerances of species present in the vegetation mosaic form the basis for variation in vegetation composition (Ross et al. 2006). Hence, the change in vegetation-inferred hydroperiod on the prairie portion of M1 and M3 reflects the amount and direction of change in vegetation (Armentano et al. 2006). In this study, the observed vegetation shift on M1 toward a drier type was not a surprise, especially considering that most years after the E1 sampling event in Spring 2006 were relatively dry. This trend will likely be reversed with the implementation of the MOD Water Delivery Project, which will send more water into the Park through NESRS.

Management-driven water conditions were also responsible for the discrepancy in vegetation change patterns observed in the eastern and western prairies on M3. Water conditions in the prairies west of SRS are influenced by the regulatory schedules for the S-12 structures along Tamiami Trail, implemented under the operational objectives of Interim Structural and Operation Plan (ISOP)/Interim Operational Plan (IOP). In concurrence with management efforts to regulate water deliveries from the S-12 structures, a consistently low water level has been maintained at water recorder NP-205. This has caused the vegetation composition to shift toward a drier type in recent years. However, such a shift towards drier vegetation was less prominent during E4 (2016) sampling, primarily because of the unusually high water conditions in the spring of 2016. These conditions were brought on by high rainfall due to the very strong El Niño, which necessitated the 2016 Temporary Emergency Deviation operated by USACE. The four-year average water depth during E4 sampling also was about 5 cm higher than the previous two samplings. The effect of spring 2016 flooding will likely be intensified and the drying trend will be reversed if the water conditions in subsequent years continue to be high. In the past, dry season flooding following by years of high water conditions have also caused significant change in habitat conditions. For instance, the management-driven spring season flooding of 1993, followed by high rains in subsequent years, caused short-hydroperiod wet prairie vegetation to change to long-hydroperiod sawgrass marsh that was less suitable habitat for CSSS (Nott et al. 1998). However, the current water management goal of moving water from west to east seems unlikely to let that happen. Recent modeling carried out using Regional Simulation Model (RSM) tool to evaluate the potential impact of Everglades Restoration Transition Plan (ERTP) also has shown that marl prairies to the northeast of current CSSS habitat will be relatively dry (USACE 2011, 2014; USFWS 2016) resulting in a major change in vegetation composition in the transition zone.

Management-driven water condition has also been a driver of the vegetation shift observed on the eastern portion of Transect M3. In this region, water pump structures at S332B and S332C deliver

water from the L31N canal into a series of inter-connected detention ponds. These ponds have a large fixed-crest weir on the western levee that allows water from the pond to enter ENP marl prairies. In addition, water may also enter ENP through subsurface flow. The purpose of operating pump stations along the L-31N canal includes lowering canal and groundwater levels and creating a continuous hydraulic ridge to control seepage back to the canal while protecting the marl prairie (sparrow habitat) from further deterioration (USACE 2006). Pumping through S332B and S332C serves the management goal of re-hydrating the marl prairies of the Rocky Glades. Thus, a shift in vegetation towards wetter types indicates that the management goal is being achieved, at least in part. However, regular monitoring is essential to detect a signal that inputs of water from the ponds continue to cause a shift in vegetation to from marl-dominated wet prairie to marsh types.

In addition to a positive outcome of the operations of water pumps and detention ponds along eastern boarder of the Park, impact of such management efforts on prairie vegetation needs to be interpreted cautiously, because water flow from detention ponds towards prairies in the Park may have adverse consequences as well. For instance, periphyton near inflow structures had elevated phosphorus in comparison to adjacent marl prairie sites to the west, suggesting an increase in P-loading due to long-term exposure of the canal-side sites to seepage (Gaiser et al. 2008; 2014). Sah et al. (2014) also concluded that vegetation in upper Taylor Slough basin showed significant trajectory along the vector representing the phosphorus gradient, possibly due to the influence of seepage water from the detention ponds. In this study, while soil TC, OC and TN increased along marl prairie-slough gradient as expected (**Figure 6**), soil phosphorous at the site near (300m) the detention pond levee was higher than at several sites 1200-2400 m from levee (**Figure 7**). If water from the detention ponds continues to influence vegetation in the adjacent prairies, the water quality issue also needs to be addressed so that the affected marl prairies do not shift to another stable state more adapted to P-enriched soil (Hagerthey et al. 2008).

In summary, regional differences in hydrologic regimes resulting from alternative management strategies have caused variation in species composition within individual landscapes and have also brought on temporal change in vegetation composition in Shark River Slough and adjacent marl prairies. The occurrence of these changes coincided with changes in hydrologic regimes during the past 17 years. Our results provide feedback for the adaptive management of Everglades wetland ecosystems along the marl prairie-slough gradient.

Acknowledgements

We would like to acknowledge the assistance in field and lab (during the period between 2005 and 2016) provided by the following members of our lab: Pablo Ruiz, Dr. Rachel King, Dave L. Reed, David Jones, Nilesh Timilsina, Mike Kline, Brooke Shamblin, Nate Colbert, Lawrence Lopez, Diana L. Rodriguez, Allison M. Lambert, Suresh Subedi, Danielle Crisostomo, Jesus Blanco, Junnio Freixa and Allison Jirout. The project received financial support from the Department of Interior's Critical Ecosystems Study Initiative (CESI), Everglades National Park (1998-2000) and the RECOVER working group within the Comprehensive Everglades Restoration Plan (CERP). The support from the RECOVER working group was provided US Army Corps of Engineers (U.S. Army Engineer Research and Development Center).

5. References

- Armentano, T. V., J. P. Sah, M. S. Ross, D. T. Jones, H. C. Cooley, C. S. Smith (2006) Rapid responses of vegetation to hydrological changes in Taylor Slough, Everglades National Park, Florida, USA. *Hydrobiologia* 569: 293-309
- Bernhardt, C. E., and D. A. Willard (2009) Response of the Everglades' ridge and slough landscape to climate variability and 20th century water-management. *Ecol. Appl.*, 19, 1723–1738.
- Busch, D. E., W. F. Loftus and O. L. Jr. Bass (1998) Long-term hydrologic effects on marsh plant community structure in the southern Everglades. *Wetlands*, 18, 230–241.
- CERP (2000) Comprehensive Everglades Restoration Plan. U.S. Army Corps of Engineer (USACE) and South Florida Water Management District (SFWMD), Florida, USA URL: <http://www.evergladesplan.org/> (last date accessed: 20 August 2012)
- EDEN (Everglades Depth Estimation Network) (2008) South Florida Information Access (Sofia). <http://sofia.usgs.gov/eden>.
- Gaiser EE, Price RM, Scinto LJ, Trexler JC (2008) Phosphorus retention and sub-surface movement through the S-332 detention basins on the eastern boundary of Everglades National Park. Year 3 Final Report to Everglades National Park, Homestead, FL, USA
- Gaiser, E. E., P. Sullivan, F. A. C. Tobias, A. J. Bramburger and J. C. Trexler (2014) Boundary effects on benthic microbial phosphorus concentrations and diatom beta diversity in a hydrologically-modified, nutrient-limited wetland. *Wetlands* 34 (Suppl 1):S55-S64.
- Hagerthey, S.E., S. Newman, K. Rutchey, E.P. Smith and J. Godin (2008) Multiple regime shifts in a subtropical peatland: community-specific thresholds to eutrophication. *Ecological Monographs* 78: 547-565.
- Juggins. S. (2003) C2 User guide. Software for ecological and palaeoecological data analysis and visualization. University of Newcastle, Newcastle upon Tyne, UK. 69pp.
- Kantvilas, G. and P. R. Minchin (1989) An analysis of epiphytic lichen communities in Tasmanian cool temperate rainforest. *Vegetatio* 84: 99-112.
- Larsen, L., N. Aumen, C. Bernhardt, V. Engel, T. Givnish, S. Hagerthey, J. Harvey, L. Leonard, P. McCormick, C. McVoy, G. Noe, M. Nungesser, K. Rutchey, F. Sklar, T. Troxler, J. Volin and D. Willard (2011) Recent and historic drivers of landscape change in the Everglades ridge, slough and tree island mosaic. *Critical Reviews in Environmental Science and Technology* 41: 344-381.
- Light, S. S. and J. W. Dineen (1994) Water control in the Everglades: a historical perspective. In: Davis S.M. and Ogden J.C. (eds), *Everglades: The Ecosystem and Its Restoration*. St. Lucie Press, Boca Raton, Florida, USA, pp. 47–84.
- McCune, B. and J. B. Grace (2002) *Analysis of Ecological Communities*. MjM Software Design, Gleneden Beach, OR, USA
- McVoy C. W., W. P. Said, J. Obeysekera, J. A. VanArman and T. W. Dreschel (2011) *Landscapes and Hydrology of the Pre-drainage Everglades*. University Press of Florida, Gainesville, FL, USA
- Minchin, P. R. (1998) DECODA: Database for Ecological Community Data. Anutech Pty. Ltd., Canberra, Australia
- Minchin, P. R., M. Folk and D. Gordon (2005) Trajectory Analysis: a New Tool for the Assessment of Success in Community Restoration. Meeting Abstract, Ecological Society of America 90th Annual Meeting, Montreal, Quebec, August 7-12, 2005

- Nungesser, M. K. (2011) Reading the landscape: temporal and spatial changes in a patterned peatland. *Wetlands Ecology and Management* 19: 475-493.
- Nott, M. P., O. L. Bass Jr., D. M. Fleming, S. E. Killeffer, N. Fraley, L. Manne, J. L. Curnutt, T. M. Brooks, R. Powell and S. L. Pimm (1998) Water levels, rapid vegetational changes, and the endangered Cape Sable seaside sparrow. *Animal Conservation* 1: 23-32
- Olmsted, I. C., and L. L. Loope (1984) Plant communities of Everglades National Park. In: P.J. Gleason, (Ed). *Environments of South Florida: Present and Past II*. pp. 167-184. Miami Geological Society, Coral Gables.
- Olmsted I. and T. V. Armentano (1997) *Vegetation of Shark Slough, Everglades National Park*. SFNRC Technical Report 97-001. South Florida Natural Resource Center, Everglades National Park, Homestead, Florida, USA
- RECOVER 2009. CERP Monitoring and Assessment Plan (MAP) – Revised 2009. Restoration and Coordination and Verification, Comprehensive Everglades Restoration Plan, Central and Southern Florida Project.
http://www.evergladesplan.org/pm/recover/recover_map_2009.aspx
- Ross, M. S., P. L. Ruiz, D. L. Reed, K. Jayachandran, J. P. Sah, and M. T. Lewin. (2001). Assessment of marsh vegetation responses to hydrological restoration in Shark Slough, Everglades National Park. Final Report submitted to Everglades National Park, Homestead, FL, USA. 70 pp.
- Ross, M. S., D. L. Reed, J. P. Sah, P. L. Ruiz, and M. T. Lewin (2003) Vegetation:environment relationships and water management in Shark Slough, Everglades National Park. *Wetland Ecology and Management* 11: 291-303.
- Ross, M. S., P. L. Ruiz, J. P. Sah, S. Stoffella, N. Timilsina, and E. Hanan (2005) Marl Prairie/Slough Gradients: Pattern and trends in Shark Slough and adjacent marl prairies. 1st Annual Report 2005.
- Ross, M. S., J. P. Sah, P. L. Ruiz, D. T. Jones, H. C. Cooley, R. Travieso, J. R. Snyder, D. Hagyar (2006) Effect of Hydrology Restoration on the Habitat of the Cape Sable Seaside Sparrow. Report to Everglades National Park, Homestead, FL, USA. 50 pp.
- Ross, M. S., J. B. Heffernan, J. P. Sah, P. L. Ruiz, A. A. Spitzig, E. Isherwood and J. Blanco. (2016). Everglades Ridge, Slough, and Tree Island Mosaics. Annual Report submitted to US Army Engineer Research and Development Center. Cooperative Agreement #: W912HZ-10-2-0030. Modification # P00004. Year 5 Report (2010-2015): 2016. 98 pp
- Ruiz, P. L., J. P. Sah, M. S. Ross and A. A. Spitzig (2013) Tree island response to fire and flooding in the short-hydroperiod marl prairie grasslands of the Florida Everglades. *Fire Ecology* 9 (1): 38 – 54.
- Sah, J. P., M. S. Ross, J. R. Snyder, P. L. Ruiz, D. T. Jones, R. Travieso, S. Stoffella, N. Timilsina, E. Hanan and H. Cooley (2007) Effect of Hydrologic Restoration on the Habitat of the Cape Sable seaside sparrow. 2005-2006. Year-4 Final Report submitted to U. S. Army Corps of Engineers, Jacksonville, FL, USA. March 2007. 49 pp.
- Sah, J. P., M. S. Ross and P. L. Ruiz. 2013. Landscape Pattern – Marl Prairie/Slough Gradient: Vegetation Composition along the Gradient and Decadal Vegetation Change Pattern in Shark Slough. Annual Report-2012 submitted to US Army Engineer Research and Development Center. Cooperative Agreement #: W912HZ-09-2-0018. Modification # P00002. April 30, 2013. 51 pp

- Sah, J. P., M. S. Ross, S. Saha, P. Minchin and J. Sadle (2014) Trajectories of vegetation response to water management in Taylor Slough, Everglades National Park, Florida. *Wetlands* (Suppl 1): S65-S79.
- U.S. Army Corps of Engineers (USACE). 2006. Interim Operational Plan for Protection of the Cape Sable seaside sparrow, Central and Southern Florida Project for Flood Control and Other Purposes, Final Supplemental Environmental Impact Statement, 2006. Jacksonville District, Jacksonville, Florida.
- U.S. Army Corps of Engineers (USACE). 2011. Everglades Restoration Transition Plan (ERTP): Final Environmental Impact Statement. U.S. Army Corps of Engineers, Jacksonville, FL.
- U.S. Army Corps of Engineers (USACE). 2014. CERP – Central Everglades Planning Project (CEPP): Final Integrated Project Implementation Report and Environmental Impact Statement. U.S. Army Corps of Engineers, Jacksonville District, FL.
- U.S. Fish and Wildlife Service (USFWS). 2016. Biological Opinion for the Everglades Restoration Transition Plan-2006. Submitted to U.S. Army Corps of Engineers, Jacksonville, FL.
- Van Lent T., R. W. Snow and F. E. James (1999) An Examination of the Modified Water Deliveries Project, the C-111 Project, and the Experimental Water Deliveries Project: Hydrologic Analyses and Effects on Endangered Species. South Florida Natural Resources Center, Everglades National Park, Homestead, Florida, USA
- Zweig, C. L. and W. M. Kitchens (2008) Effects of landscape gradients on wetland vegetation communities: information for large-scale restoration. *Wetlands* 28:1086-1096
- Zweig, C. L., and W. M. Kitchens (2009). Multi-state succession in wetlands: A novel use of state and transition models. *Ecology*, 90, 1900–1909.

Appendix 1: Importance value index (IV) of species present at the marl prairie sites of Transect M1, M2E, M3E and M3W that were sampled four times between 2006 and 2014. M3E and M3W are the part of M3 on east and west of Shark River Slough, respectively.

S.NO	Species	M3E													
		M1				M2E	M3E				M3W				
		2006	2009	2012	2015	2015	2007	2010	2013	2016	2007	2010	2013	2016	
1	<i>Aeschynomene pratensis</i>	0.81	1.52	0.06	2.19	0.49			0.01		0.12	0.79	0.56	0.09	1.29
2	<i>Agalinis linifolia</i>		0.46						0.20	0.02	0.27	0.15	0.07	0.11	0.87
3	<i>Agalinis</i> sp.									0.00					
4	<i>Aletris bracteata</i>								0.02	0.01	0.05				
5	<i>Ambrosia artemisiifolia</i>										0.04				
6	<i>Andropogon glomeratus</i> var. <i>glomeratus</i>							0.01			0.05				
7	<i>Andropogon virginicus</i>	0.69	1.05	2.43	0.24	0.14	0.02	0.46	0.37	0.26	0.34	0.46	0.02	0.23	
8	<i>Anemia adiantifolia</i>						0.11								
9	<i>Angadenia berteroi</i>						0.10	0.05	0.00	0.08					
10	<i>Annona glabra</i>	0.42		0.18	0.15	0.45	0.08	0.06	0.06	0.06	3.58	1.96	0.60	0.78	
11	<i>Ardisia escallonioides</i>							0.01	0.00	0.02					
12	<i>Aristida purpurascens</i>			0.38			0.04	0.95	0.74	0.94	0.02	0.13	0.56	0.29	
13	<i>Aristida stricta</i>						0.02								
14	<i>Asclepias lanceolata</i>					0.03	0.16	0.07	0.00	0.04	0.03	0.16	0.08	0.23	
15	<i>Asclepias longifolia</i>												0.04	0.04	
16	<i>Symphyotrichum adnatum</i>					0.07									
17	<i>Symphyotrichum bracei</i>				2.94	0.92				2.05	0.33			1.19	
18	<i>Symphyotrichum dumosum</i>				0.47		0.01	0.08	0.02	0.38	0.64	2.23	0.32	1.04	
19	<i>Symphyotrichum subulatum</i>						0.92								
20	<i>Symphyotrichum tenuifolium</i>	1.75	1.17	3.09				4.99	0.66	0.11	0.06	2.54	0.59	0.36	
21	<i>Aster</i> sp.						0.08								
22	<i>Bacopa caroliniana</i>	3.94	1.14	1.50	4.95	4.65	2.27	1.95	0.95	1.21	4.64	4.43	1.37	3.11	
23	<i>Baccharis glomeruliflora</i>									0.13					
24	<i>Baccharis halimifolia</i>		0.04				0.11		0.08						
25	<i>Bacopa monnieri</i>										0.04	0.48			
26	<i>Blechnum serrulatum</i>					0.07	0.59	0.10	0.30	0.27					
27	<i>Buchnera americana</i>						0.01			0.04				0.11	

S.NO	Species	M3E												
		M1				M2E	M3E				M3W			
		2006	2009	2012	2015	2015	2007	2010	2013	2016	2007	2010	2013	2016
28	<i>Capraria biflora</i>								0.00					
29	<i>Cassytha filiformis</i>						0.70	0.29	0.05	0.19	3.81	0.59	0.71	1.37
30	<i>Centella asiatica</i>	2.77	3.00	3.56	4.63	1.54	3.27	0.70	0.40	0.16	3.04	4.64	2.49	2.85
31	<i>Cephalanthus occidentalis</i>				0.17				0.00		0.01		0.00	0.01
32	<i>Chiococca parvifolia</i>					0.29	0.04			0.05				
33	<i>Chrysobalanus icaco</i>									0.07				
34	<i>Cirsium horridulum</i>							0.05	0.00	0.01				
35	<i>Cladium mariscus ssp. jamaicense</i>	40.94	41.35	37.42	27.65	34.49	47.57	38.03	22.79	42.34	20.38	22.39	24.44	25.86
36	<i>Coelorachis rugosa</i>											0.01		
37	<i>Conoclinium coelestinum</i>							0.42						
38	<i>Crinum americanum</i>		0.31	0.54	0.33	0.23				0.01	1.72	1.03	0.53	2.37
39	<i>Cyperus haspan</i>			0.06			0.01	0.14	0.01					
40	<i>Cyperus sp.</i>							0.08						
41	<i>Dichantherium aciculare</i>			0.04		0.11			0.03	0.15				
42	<i>Dichantherium dichotomum</i>				0.16	0.06	1.11	1.64	0.01	1.76				0.13
43	<i>Dichantherium sp.</i>					0.33								
44	<i>Diodia virginiana</i>								0.00					
45	<i>Dyschoriste angusta</i>					0.11							0.07	
46	<i>Echinochloa sp.</i>										0.01			
47	<i>Eleocharis baldwinii</i>			0.04					0.03					
48	<i>Eleocharis cellulosa</i>	9.88	13.72	9.91	7.14	27.15	3.25	4.94	3.59	3.59	2.35	2.08	1.28	2.43
49	<i>Eleocharis elongata</i>							0.08						
50	<i>Eleocharis geniculata</i>				0.06									
51	<i>Eragrostis elliottii</i>	0.27		0.84	0.16	0.07	0.87	0.81	0.03	0.32	0.20	0.32	0.15	0.37
52	<i>Eriocaulon compressum</i>											0.15		0.04
53	<i>Erigeron quercifolius</i>						0.07	0.12	0.01		0.06			
54	<i>Eugenia axillaris</i>						0.08		0.00					
55	<i>Eupatorium capillifolium</i>		0.36	0.04			1.30							
56	<i>Eupatorium leptophyllum</i>				0.26			0.49	0.16	0.37				
57	<i>Eupatorium mikanioides</i>		0.29	0.04	0.21		0.18	0.11	0.00	0.02	0.16	0.24	0.08	0.53

S.NO	Species	M3E												
		M1				M2E	M3E				M3W			
		2006	2009	2012	2015	2015	2007	2010	2013	2016	2007	2010	2013	2016
58	<i>Eustachys petraea</i>						0.11		0.12				0.04	
59	<i>Flaveria linearis</i>						0.36							
60	<i>Fuirena breviseta</i>	0.07		0.21	0.03	0.06	0.01	0.59	0.15	0.12	0.01		0.07	
61	<i>Habenaria repens</i>											0.07		
62	<i>Helenium pinnatifidum</i>											0.01	0.11	
63	<i>Heliotropium polyphyllum</i>					0.20	0.15	0.09	0.00	0.11				
64	<i>Hibiscus grandiflorus</i>						0.01			0.05	0.07	0.01	0.08	
65	<i>Hydrolea corymbosa</i>								0.00					
66	<i>Hymenocallis latifolia</i>												0.66	
67	<i>Hymenocallis palmeri</i>						0.17	0.07	0.05	0.41	1.16	1.18	0.62	
68	<i>Hyptis alata</i>						0.38	0.25	0.04	0.27	0.04	0.10	0.05	
69	<i>Hypericum cistifolium</i>							0.07						
70	<i>Hypericum hypericoides</i>								0.00					
71	<i>Ipomoea sagittata</i>		1.20	0.03	0.71	0.48	0.27	0.15	0.16	0.62	0.37	0.34	0.10	
72	<i>Iva microcephala</i>			0.59	0.77	0.40	0.14	0.46	0.08	0.12				
73	<i>Juncus megacephalus</i>												0.01	
74	<i>Justicia angusta</i>	0.90	0.96	1.33	1.44	0.47	0.27	1.02	0.08	1.16	0.96	1.56	0.47	
75	<i>Kosteletzkya virginica</i>							0.03	0.10				0.02	
76	<i>Leersia hexandra</i>	0.21	0.78	0.35	0.19			0.16			0.23	0.70	0.31	
77	<i>Linum medium</i>						0.01	0.02					0.03	
78	<i>Lobelia glandulosa</i>							0.03	0.01		0.14		0.16	
79	<i>Ludwigia alata</i>				0.33	0.06	0.03	0.34	0.10	0.12			0.00	
80	<i>Ludwigia curtissii</i>				0.30	0.03				0.36				
81	<i>Ludwigia microcarpa</i>	0.22	0.06	0.93	0.04		0.19	0.63	0.11	0.10	0.06	0.80	0.07	
82	<i>Ludwigia repens</i>		1.38			0.06	1.20	0.81	0.08		0.32	0.05		
83	<i>Ludwigia sp.</i>									0.44				
84	<i>Melaleuca quinquenervia</i>		0.22											
85	<i>Metopium toxiferum</i>						0.20							
86	<i>Mikania scandens</i>	0.27		0.18		0.07	1.55	1.84	0.09	1.06	0.14	0.14	0.05	
87	<i>Mitreola petiolata</i>			0.23		0.20	0.71	0.08	0.00	0.39	0.81	0.28	0.18	

S.NO	Species	M3E												
		M1			M2E		M3E				M3W			
		2006	2009	2012	2015	2015	2007	2010	2013	2016	2007	2010	2013	2016
88	<i>Muhlenbergia capillaris</i> var. <i>filipes</i>	2.46	5.42	4.32	7.90	3.17	7.49	7.74	3.55	7.07	0.61	0.39	0.18	0.17
89	<i>Morella cerifera</i>	0.12			0.07	0.15	0.62	0.01	0.30	0.67	0.06	0.17	0.03	0.09
90	<i>Myrsine floridana</i>						0.14			0.05				
91	<i>Nymphoides aquatica</i>						0.01	0.03			0.07	0.20	0.00	0.18
92	<i>Nymphaea odorata</i>											0.05		
93	<i>Oxypolis filiformis</i>	0.50	0.14	0.09	0.09	0.04	0.11	0.48	0.04	0.10	0.16	0.18	0.17	0.21
94	<i>Panicum dichotomiflorum</i>							1.57						
95	<i>Panicum hemitomom</i>	0.21		0.07	1.16	1.38	0.22	0.93	0.20	0.14	2.15	1.54	0.20	0.60
96	<i>Panicum rigidulum</i>						0.04	0.13	0.18		0.04	0.14	0.00	
97	<i>Panicum tenerum</i>	4.38	4.35	5.74	3.65	1.45	3.05	5.43	2.22	5.06	3.37	6.77	1.75	4.95
98	<i>Panicum virgatum</i>		0.61	0.03	0.33	0.41	0.61	0.15	0.06	0.59	5.59	7.57	5.06	6.58
99	<i>Panicum</i> sp.							0.11						
100	<i>Parthenocissus quinquefolia</i>								0.00	0.15				
101	<i>Paspalidium geminatum</i>				0.17	1.55	0.05		0.11	0.12	0.14	0.61	0.02	0.28
102	<i>Paspalum monostachyum</i>							0.05	0.06	0.21	4.29	8.47	5.60	5.02
103	<i>Passiflora suberosa</i>							0.01	0.02					
104	<i>Peltandra virginica</i>	0.74	1.16	0.21	0.06		0.11	0.06	0.01	0.27	0.01		0.12	0.06
105	<i>Persea borbonia</i>						0.20	0.10	0.38	0.75	0.03	0.13	0.00	0.03
106	<i>Phytolacca americana</i>								0.01					
107	<i>Phyllanthus caroliniensis</i>					0.05	0.05							
108	<i>Phyla nodiflora</i>	1.39	0.51	0.44	2.66	1.08	3.77	0.89	0.58	0.60	0.10	0.09		
109	<i>Phyla stoechadifolia</i>		0.74	0.47	0.89	0.60	0.70	0.10	0.04	0.32				
110	<i>Phyllanthus</i> sp.												0.02	
111	<i>Pinguicula pumila</i>						0.01							
112	<i>Piriqueta cistoides</i> ssp. <i>caroliniana</i>												0.02	
113	<i>Pluchea odorata</i>									0.03				
114	<i>Pluchea rosea</i>	1.61	2.67	1.55	2.78	1.09	4.10	4.50	1.02	4.51	3.40	3.20	1.00	2.22
115	<i>Polygala baldunii</i>					0.07	0.01							
116	<i>Polygala grandiflora</i>						0.10	0.04	0.00	0.04			0.03	0.13
117	<i>Polygonum hydropiperoides</i>		0.79	0.75	0.06	0.19	0.31	0.15	0.06	0.09	0.17	0.09	0.02	

S.NO	Species	M3E												
		M1				M2E	M3E				M3W			
		2006	2009	2012	2015	2015	2007	2010	2013	2016	2007	2010	2013	2016
118	<i>Pontederia cordata</i>	0.27	0.58	0.27	0.33		0.35	0.00		0.42	0.41	0.02	0.06	
119	<i>Proserpinaca palustris</i>		0.72	0.19	0.07		0.75	0.54	0.19	0.24	0.30	0.06	0.16	
120	<i>Psychotria nervosa</i>						0.35			0.01				
121	<i>Pteridium aquilinum</i>						0.58	0.34	0.25	0.20				
122	<i>Quercus</i> sp.												0.01	
123	<i>Randia aculeata</i>						0.38	0.01	0.08	0.26				
124	<i>Rhus copallinum</i>								0.25	0.09				
125	<i>Rhynchospora colorata</i>						0.01	0.15	0.03	0.01	0.13	0.06	0.07	
126	<i>Rhynchospora divergens</i>				0.03	1.11		0.85	0.86	0.01			0.09	
127	<i>Rhynchospora inundata</i>	0.25		1.12	1.35	0.65	0.12			0.09	4.66	1.70	1.31	
128	<i>Rhynchospora microcarpa</i>	0.05		1.35	0.44	1.76	0.69	1.38	0.71	1.81	9.21	3.00	3.16	
129	<i>Rhynchospora miliacea</i>		1.56											
130	<i>Rhynchospora tracyi</i>	11.80	9.08	13.12	15.89	9.57	0.61	4.54	1.08	2.70	10.18	4.70	5.77	
131	<i>Rhynchospora</i> sp.						0.02							
132	<i>Sabatia grandiflora</i>						0.06	0.14		0.04	0.12		0.29	
133	<i>Sabal palmetto</i>					0.03	0.01	0.02			0.01			
134	<i>Saccharum giganteum</i>		0.18	0.22	0.60		0.18	0.06	0.01	0.21	0.01	0.07	0.18	
135	<i>Sagittaria lancifolia</i>				0.15	0.16	0.61	0.59	0.12	0.53	0.77	1.04	0.58	
136	<i>Salix caroliniana</i>	0.21	0.22	0.32	0.33		0.08		0.28	0.03	0.12		0.36	
137	<i>Samolus ebracteatus</i>					0.05	0.37	0.23	0.05	0.16				
138	<i>Funastrum clausum</i>										0.18	0.38		
139	<i>Schoenolirion albiflorum</i>											0.19	0.05	
140	<i>Schoenus nigricans</i>						0.01	0.01	0.05	0.01	1.24	1.73	2.21	
141	<i>Schizachyrium rhizomatum</i>		0.15	0.24		0.11	2.97	2.72	1.85	5.32	4.64	5.71	8.78	
142	<i>Schinus terebinthifolius</i>					0.01	0.08		0.01					
143	<i>Scirpus</i> sp.										0.09			
144	<i>Setaria parviflora</i>			0.69	1.58	0.37	0.74	0.08		0.65	0.01	0.22		
145	<i>Sideroxylon salicifolium</i>						0.01	0.05	0.00					
146	<i>Smilax laurifolia</i>						0.02							
147	<i>Solidago fistulosa</i>							0.16						

S.NO	Species	M3E												
		M1				M2E	M3E						M3W	
		2006	2009	2012	2015	2015	2007	2010	2013	2016	2007	2010	2013	2016
148	<i>Solidago stricta</i>		0.03	0.13		0.39	0.29	0.82	0.08	0.25	0.17		0.02	0.18
149	<i>Taxodium distichum var. imbricrium</i>										0.05			
150	<i>Teucrium canadense</i>					0.01		0.29	0.04	0.04		0.11		
151	<i>Thalia geniculata</i>							0.03			0.01	0.12	0.02	
152	<i>Thelypteris kunthii</i>						0.01							
153	<i>Toxicodendron radicans</i>						0.39	0.06	0.21	0.65				
154	<i>Trema micrantha</i>							0.22	0.07	0.02				
155	<i>Typha domingensis</i>	6.34	0.94	1.84	3.82						0.35	0.67	1.17	0.38
156	<i>Unknown grass</i>									0.05	0.03			0.05
159	<i>Unknown sp01</i>										0.06			
160	<i>Unknown sp02</i>								0.00		0.14			
161	<i>Unknown sp03</i>								0.00		0.04			
162	<i>Unknown sp04</i>										0.08			
163	<i>Unknown sp05</i>										0.05			
164	<i>Unknown sp06</i>							0.10						
165	<i>Unknown sp07</i>											0.04		
166	<i>Unknown sp08</i>										0.05			
157	<i>Unknown sp16</i>					0.21								
158	<i>Unknown sp17</i>									0.02				
167	<i>Utricularia cornuta</i>					0.34							0.02	
168	<i>Utricularia foliosa</i>	0.05	0.09	0.51			0.08	0.01	0.00	0.96		0.02		0.56
169	<i>Utricularia purpurea</i>	6.49	1.04	2.36				0.09	1.43	4.32		0.42		1.28
170	<i>Utricularia subulata</i>											0.01		
171	<i>Utricularia sp.</i>							0.01						
172	<i>Vernonia blodgettii</i>							0.16	0.00	0.02				
173	<i>Vitis rotundifolia</i>						0.16			0.04				